





Kiama Municipal Council


Annual Surface and Groundwater Monitoring Report (EPL) Gerroa Waste Disposal Depot

Report E2W-025 R001

11 March 2016



Prepared by: D.Parisotto(Managing Director) BAppSc- Geology (Hons);MAppSc- Groundwater Phone: (02) 4234 0829 Fax: (02) 4236 1824 175 Fern Street, Gerringong,NSW,Australia,2533







Client: Kiama Municipal Council

Project: Annual Surface and Groundwater Monitoring Report Gerroa Waste Disposal Depot

> Prepared for: Paul Czulowski Kiama Municipal Council P.O. Box 75, Kiama, NSW, 2533.

> Report: 11 March 2016 Reference: E2W-025 Gerroa R001 (V1)

Water Sampling & Laboratory Analyses Completed By: ALS Environmental Pty Ltd

Authorised by: Earth2Water Pty Ltd

Dino Parisotto - Director

BAppSc Geology (Hons); MAppSc (Groundwater)

Ph: (02) 4234 0829 Fax: (02) 4236 1824 175 Fern Street, Gerringong, NSW, 2533

Reports Distributed to:

Kiama Municipal Council: pdf copy

© Authorisation from Earth2Water Pty Ltd is required for third party use and distribution



TABLE OF CONTENTS

1	Introduction	. 4
	1.1 Background and Remediation activities1.2 objectives	
2	Scope of Work	. 5
3	Licence Criteria and Relevant Guidelines	. 6
4	Environmental Setting	. 8
	4.1 Climate	. 8 . 9 . 9
5	Previous Monitoring Results	11
6	Surface and Groundwater monitoring	11
	6.1 Monitoring locations	12 13
7	Monitoring Results	14
	7.1 Groundwater data	14 15 15 16 16 16 18 19 20 20 20 21 21 21
	7.2.2.2 Total Phosphorous (TP)	23
	7.2.3 Bacteriological Contaminants	



		Major Ions Quality Assurance/Quality Control	
8	Lead	chate plume and Landfill Rehabilitation	25
	8.1	Ecological Issues	26
9	Con	clusions	26
	9.1	Recommendations	28
10	Lim	itations	29
11	Refe	erences	30

TABLES

Table GW-1: Summary Analytical Report - Groundwater

Table SW-1: Summary Analytical Report - Surface Water

Table 6: Groundwater and Surface Water Monitoring (May 2015 to February 2016)

GRAPHS

Graph-01: Groundwater Ammonia Time-Series Trend - Standard & Creek Wells

Graph-02: Groundwater Ammonia Time-Series Trends - Deep Wells

Graph-03: Groundwater Total Phosphorous Time Series Trends -Standard wells

Graph-04: Groundwater Total Phosphorous Time-Series Trends - New Shallow & Deep Wells

Graph-05: Depth to Groundwater (m AHD) 2001 to February 2016

Graph-06: Surface Water Ammonia Time-Series Trends

Graph-07: Surface Water Time Series - Total Phosphorous Trends

FIGURES

Figure 1: Site Location

Figure 2: Site Layout & Well Locations

Figure 3A: Inferred Groundwater Flow Regime (Dry, August 2015) Figure 3B: Inferred Groundwater Flow Regime (Wet, May 2015)

Figure 4: Ammonia Results mg/L (2015-2016)

APPENDICES

Appendix A: Laboratory Certificates and Field Records

Appendix B: Rainfall Data

Appendix C: Ammonia Trigger Values



1 INTRODUCTION

Earth2Water Pty Ltd (E2W) was engaged by Kiama Municipal Council (Council) to provide an annual groundwater and surface water monitoring report for the Gerroa Waste Disposal Depot (GWDD). The GWDD Environment Protection Licence (EPL) was revoked in May 2008, and altered the previous reporting periods from August-July, to 1 April 2008 - 31 March 2009. The EPL reporting periods and monitoring periods have changed since 2009 (i.e. 2009-2010 monitoring report included results from five monitoring events including February 2009, May 2009, August 2009, November 2009 and February 2010. The 2010-2011 monitoring report included results from four monitoring events including May 2010, August 2010, November 2010 and February 2011. The 2011-2012 annual report includes results from five monitoring events including May 2011, August 2011, November 2011, February 2012 and May 2012. The 2012-2013 and 2013-2014/2014-2015 annual reports include results from four regular monitoring events (May, August, November and February).

This annual EPL report (2015-2016) by E2W includes an assessment of four monitoring events (May 2015, August 2015, November 2015 and February 2016) and is based on Council's project brief (Ref H24/11, August 2004), ALS Environmental Pty Ltd laboratory results and E2W previous reports. This EPL report (2015-2016) is the twelth provided by E2W to the NSWEPA on behalf of Council, and meets the conditions outlined in the EPL (Lic No: 5959, R1.10).

1.1 BACKGROUND AND REMEDIATION ACTIVITIES

Council has owned and operated the GWDD since the 1960's. It was previously licensed as a Solid Waste Class 1 Landfill, operating under the EPL No. 5959. The site also functioned as a night soil depot for liquid pump out sullage. Land filling operations at the GWDD were discontinued in October 2003.

From July 2004 to February 2005, the landfill was rehabilitated to eliminate, or at least minimise the potential for landfill leachate generation. The waste mound was reshaped, capped with a 0.5 m thick clay barrier and 0.3 m thick combined drainage/re-vegetation layer. The former night soil trench was also remediated in August 2004 (i.e. approximately 300 tonnes of bio-solid sludge were excavated and placed underneath the clay cap).

In February 2005, an irrigation system was installed to assist with the re-vegetation of the landfill mound using a combination of native and annual grasses. A groundwater holding dam (30 x 30 m) was also constructed next to the two existing evaporation ponds to contain and supply water for the irrigation system. A spear point (yield \sim 2 L/sec) was installed on the north-west corner of the landfill mound to allow groundwater (and landfill leachate) to fill the holding dam as well as supply water for the irrigation system.

The sludge pond (southern lined dam) at Gerroa Landfill has not been used since Council ceased undertaking the septic clean-outs. Waste Processing Solutions Pty Ltd was engaged by KMC in September 2009 to de-water the sludge in the lined pond, and subsequently taken to a Soilco Pty Ltd owned site. Removal of the pond liner (HDPE) was undertaken in October 2011 by Council and

.

¹ First annual report for 2003 - 2004 was submitted October 2004.



disposed to Shellharbour landfill (note: spillage of residual sludge may have occurred during removal of the liner).

Up until November 2008, Ecowise Pty Ltd (now ALS) performed the quarterly surface and groundwater monitoring at the landfill site. E2W and Council undertook the monitoring in November 2008, and subsequently Council and/or ALS performing the quarterly monitoring rounds herein. Water samples are sent to ALS for laboratory analyses. The landfill is kept locked at all times.

1.2 **OBJECTIVES**

Similar to previous years, the objective of monitoring was to assess the potential impact of the GWDD on local surface and groundwater systems. This round of monitoring provides an assessment of water quality improvements associated with the landfill rehabilitation works completed in February 2005.

2 SCOPE OF WORK

E2W was commissioned by Council to collate and interpret surface and groundwater data from the GWDD on 27 May, 24 August, 4 November 2015, and 16 February 2016. Each monitoring event comprised the following:

- Sampling of onsite and offsite groundwater wells MW-1S, MW-1D, MW-3, MW-4, MW-5, MW-6S, MW-6D, MW-7S, MW-7D, MW-9, MW-10 and MW-11.
- Sampling of surface water at two locations along Blue Angle Creek (ML-2, and ML-5²). Only ML-2 and ML-5 locations were sampled during this reporting period (i.e. ML-1, ML-3 and ML-4 were inaccessible due to being located on private property belonging to Cleary Brothers).

Similar EPL reports were previously completed by E2W for the 2003-2004, 2004-2005, 2005-2006, 2006-2007, 2007-2008, 2008-2009, 2009-2010, 2010-2011, 2011-2012, 2012-2013, 2013-2014 and 2014-2015 monitoring periods. E2W is required to conduct the following scope of work to satisfy Council's surface and groundwater monitoring program at GWDD:

- Assist Council to interpret quarterly sampling results and provide recommendations.
- Prepare this annual report for May 2015 to February 2016 to provide information in accordance with Section R1.10 of the EPL (No. 5959). The annual report is to include the following:
 - 1. Tabulation of the monitoring data obtained for the period.
 - 2. Graphical representation of the current and previous monitoring data (minimum last three years). Statistically significant variations or anomalies will be highlighted.
 - 3. Analyses and interpretation of monitoring data.
 - 4. Analyses and response to any complaints received.
 - 5. Identification of any deficiencies in the environmental performance of the GWDD, as highlighted by the monitoring data, trends and/or accidents.
 - 6. Proposal of recommendations to address the above identified deficiencies.
 - 7. Recommendations on improving the overall environmental performance of the facility.

e2W

² November 2008 was the first time ML-5 had been sampled since October 2004.



3 LICENCE CRITERIA AND RELEVANT GUIDELINES

The EPL for the GWDD was revoked by the DECC in May 2008. The ongoing groundwater monitoring is undertaken to assess the effectiveness of the capping works and environmental status of the landfill post closure and rehabilitation. The revocation notice is subject to the following conditions:

- The licensee must maintain the landfill capping works.
- The licensee must undertake groundwater monitoring at groundwater wells MW-1S, MW-1D, MW-3, MW-4, MW-5, MW-6S, MW-6D, MW-7S, MW-7D, MW-9, MW-10 and MW-11 (in accordance with Table 3.1 below).
- Should the monitoring results indicate ammonia concentrations greater than 20% above ammonia concentrations reported in Table GW-1 of *Kiama Municipal Council, Gerroa Waste Disposal Depot Annual Groundwater and Surface Water Monitoring Report August 2006 to May 2007*, dated 17 August 2007, the licensee must notify the EPA within 7 days of receiving the results.
- The licensee must undertake surface water monitoring at surface water monitoring points ML-1, ML-2, ML-3, ML-4 and ML-5 (in accordance with Table 3.1 below).
- Should the monitoring results indicate ammonia concentrations greater than 10% above ammonia concentrations reported in Table SW-1 of *Kiama Municipal Council, Gerroa Waste Disposal Depot Annual Groundwater and Surface Water Monitoring Report August 2006 to May 2007*, dated 17 August 2007, the licensee must notify the EPA within 7 days of receiving the results (Appendix C).
- The licensee must dewater, clean out and cap the two HDPE lined liquid storage ponds
 - o By 1 January 2009, or
 - o When the ponds are no longer required, or
 - o If the liner is breached (whichever is the earlier).

Table 3.1: Surface and Groundwater Monitoring Requirements

Parameters	Monitoring Frequency - Groundwater	Monitoring Frequency – Surface water
Alkalinity	Quarterly (#1)	Quarterly (#1)
pН	Quarterly (#1)	Quarterly (#1)
Conductivity	Quarterly (#1)	Quarterly (#1)
Total Dissolved Solids	Quarterly (#1)	Quarterly (#1)
Nitrogen (Ammonia)	Quarterly (#1)	Quarterly (#1)
Phosphorous (Total)	Quarterly (#1)	Quarterly (#1)
Nitrate	Quarterly (#1)	Quarterly (#1)
Nitrite	Quarterly (#1)	Quarterly (#1)
Total Kjeldahl Nitrogen	Quarterly (#1)	Quarterly (#1)
Calcium	Annual	Annual
Chloride	Annual	-
Fluoride	Annual	-
Magnesium	Annual	Annual
Sulphate	Annual	-
Sodium	Annual	Annual
Bicarbonate	Annual	Annual
Carbonate	Annual	Annual
Potassium	Annual	Annual



Dissolved Organic Carbon	Annual	-
Iron	Annual	Annual
Manganese	Annual	Annual
Redox Potential	Annual	Annual
Faecal Coliforms	-	Annual
Enterococci	-	Annual

Note: Bicarbonate/alkalinity was conducted on quarterly basis (only an annual requirement). Quarterly (#1) = monitoring is conducted on quarterly basis by KMC, however EPL requires sampling only "twice a year".

The parameters listed in Table 3.1 were included in May, August, November 2015 and/or February 2016 (Table 6). Annual parameters were sampled during the August 2015 monitoring period. Ongoing monitoring with the same parameters is proposed for the future monitoring reports.

The DECC's Contaminated Sites: Guidelines for Assessment and Management of Groundwater Contamination (March 2007) outlines a best-practice framework for assessing and managing groundwater contamination. The guidelines are made under the Contaminated Land Management Act (1997) and recommend adopting the ANZECC (2000) chemical concentration trigger values for the protection of (fresh and marine) aquatic ecosystems.

The Australian and New Zealand Guidelines for Fresh and Marine Water Quality (ANZECC, 2000) guidelines include risk-based trigger levels and indicative interim working levels (IIWLs). The IIWLs are of low reliability and used when insufficient data is available to calculate a trigger level. It should be noted that the ANZECC (2000) water quality guidelines are applicable to receiving water and not to groundwater. However, they form an appropriate basis for undertaking a screening level assessment of groundwater quality. The selection of the applicable guideline values should be based on an assessment of potential pathways by which human or environmental exposure might take place and the beneficial end use of the groundwater (i.e. ecosystem support).

The choice of a beneficial use classification for groundwater at the site depends on the quality of the water and its potential use in the long term. Although groundwater in the aquifer surrounding the waste disposal facility is likely to be of relatively good quality (depending on the presence of saline intrusions), there are no known groundwater extraction bores (agricultural or domestic) within the immediate vicinity. Therefore, the most appropriate beneficial use category of the groundwater is considered to be for the protection of aquatic ecosystems in the discharge zones of nearby Blue Angle Creek and Seven Mile Beach (fresh and marine water, respectively).

Based on the closest environmental receptors being both marine and fresh waters (i.e. Seven Mile Beach, Blue Angle Creek and Crooked River Estuary), the guidelines adopted for the site are based on the protection of both marine and fresh water aquatic ecosystems. These assessment guidelines are presented with the summarised analytical results (i.e. Tables GW-1, SW-1).

Exceedances of ANZECC (2000) trigger values for marine water ecosystems have been highlighted in Tables GW-1 and SW-1.



4 ENVIRONMENTAL SETTING

The GWDD is located approximately 1.5 km southwest of the Gerroa Road bridge crossing of Crooked River and near the northern end of Seven Mile Beach. Blue Angle Creek is located around 80 m to the northwest of Crooked River Road (Figure 1).

The facility covers an area of approximately 3.2 hectares. The location and general layout of the site is shown in Figures 1 and 2. Prior to the remediation of the site in February 2005, the GWDD comprised the following:

- An elevated landfill mound ranging from 4 m to ~15 m AHD,
- Two lined evaporation sludge ponds. These accepted septic sludge associated with Gerringong's upgraded sewerage system. These ponds are now lined and used to hold groundwater pumped from the site to enable irrigation on the landfill mound.
- A night soil deposit, which historically accepted night soil sludge; and
- A small recycling facility.

The former night soil depot was located adjacent to the north-western slope of the landfill mound (Figure 2). The former depot consisted of two excavated infiltration trenches which, up until August 2002, received pump-out wastewater from septics in the Gerringong/Gerroa region. The two trenches were approximately 100 m in length (5 m wide) and while operational, partially filled with untreated wastewater.

4.1 CLIMATE

Between 1895 and 2011, the average yearly rainfall at Gerringong Mayflower Village (Latitude 34.75° S, Longitude 150.82° E) is 1343.4 mm/year, with the heaviest rainfalls occurring in summer and autumn months.

However, rainfall data from Toolijooa (Nyora) Station is taken from July 2012 due to the closure of Mayflower Village Station. Climatic data indicates Gerringong/Gerroa received approximately 1559 mm of rain from February 2015 to February 2016³ (Appendix B). The highest months of above average rainfall were recorded in April 2015 (426.4 mm), and August 2015 (378 mm), whilst January 2015 (316 mm i.e. just prior to reporting period).

4.2 TOPOGRAPHY

A general layout of the site topography is presented in Figure 1. The GWDD is located in an estuarine landscape consisting of dune ridges, swamps and lagoons. The vegetation surrounding the facility comprises scrub and a littoral rainforest. Local relief is less than 5 m AHD and slopes less than 5%. The landfill forms a mound, reaching ~12 m above the surrounding ground surface.

The reduced groundwater level (RL) of the landfill footprint area ranges from \sim 3.3 to 5.2 m AHD. The footprint area (23,000 m²) and height of the rehabilitated landfill mound (15.9 m AHD, July 2005) is practically the same as pre-remediation conditions (i.e. October 2003).

,

³ Information obtained from the Bureau of Meteorology website (www.bom.gov.au/climate).



4.3 GEOLOGY

The GWDD is located within the Seven Mile Soil Landscape as defined by Hazelton (1992). Coastal Plain Quaternary marine sands and Quaternary alluvium underlie the site.

Previous site investigations, as discussed in the URS report (2002), identified light brown fine-grained dune sands to a depth of 4 - 6 m below grade. The sands vary in thickness (between dune ridges and swales) and are largely contained above the groundwater table.

Beneath the dune sands, fine-grained grey beach sands containing shell fragments are present to a depth of approximately 14 m. The base of the sequence comprises estuarine silty and sandy clays. Further inland, the sedimentary sequence comprises estuarine clays closer to the surface, particularly in the vicinity of Blue Angle Creek where the ground surface is 2 - 3 m lower than the landfill area.

4.4 HYDROGEOLOGY

The GWDD is underlain by an unconfined and permeable sandy aquifer. Groundwater is encountered at a depth of approximately 3 - 4 m below ground level (i.e. ~1 m AHD) at the landfill area and becomes shallower towards Blue Angle Creek to the west and Seven Mile Beach to the east.

The groundwater quality varies from potable to saline, with electrical conductivities ranging from 0.3 to 60 mS/cm. The groundwater generally becomes more saline with depth and in the vicinity of tidal saline water bodies (i.e. Blue Angle Creek and Seven Mile Beach).

The hydraulic conductivity of the beach and dune sands at Gerroa has a geometric average of 10 m/day (Gerroa-Gerringong Sewerage Scheme EIS, 1999). It is noted that the organic silty sands between 0 - 0.15 mbgl have a lower permeability (approximately 0.1 m/day, E2W site investigations, March 2004).

In the area of the GWDD, groundwater gradients are controlled by topography, the Seven Mile Beach shoreline, Blue Angle Creek and Crooked River Estuary. It is interpreted that a natural groundwater divide runs through the landfill (URS 2003, E2W 2004), with groundwater to the west flowing towards Blue Angle Creek and groundwater to the east towards Seven Mile Beach. It is likely the natural groundwater divide is influenced by the coastal dunes and presence of the landfill mound (due to increased recharge). The position of the groundwater divide may change with the tide and seasons. During 2004/2005 (a drought period), the predominant flow direction was considered to be east towards Seven Mile Beach.

Groundwater discharge at Blue Angle Creek and Seven Mile Beach will be influenced by the presence of a fresh groundwater/salt water interface. The interface results from the density difference between the groundwater and sea water and is a dynamic and complex region with upward hydraulic gradients, tidal fluctuations, micro-biological processes, groundwater and surface interaction and substantial salinity variations. The groundwater/salt water interface can be associated with enhanced natural attenuation (biodegradation, dilution, sorption etc.), which acts to reduce the levels of contaminants prior to their discharge to marine ecosystems.



Groundwater is also interpreted to discharge as baseflow within Blue Angle Creek. Due to the action of tides, salt water is intermittently present in Blue Angle Creek with salinity governed by tide levels. The saline water intrusion at high tide extends approximately 2 km upstream of the confluence between Blue Angle Creek and Crooked River.

It is understood offsite migration of contaminants (nutrients, iron and some ammonia) has occurred in the local shallow and deep groundwater systems (URS 2002, 2003; E2W 2004, 2006). This groundwater contamination arises from nutrient enrichment, which is associated with the former landfilling operations at the GWDD.

The leachate plume identified in the well network arises from historical waste disposal at the site, which commenced in the 1960s. The landfill remediation (completed in February 2005) will reduce future landfill leachate generation, however shrinking/diminishing of the existing plume will depend on natural attenuation processes (i.e. dilution, adsorption, biodegradation dispersion etc.) over time (years).

The groundwater quality immediately outside the footprint area is subjected to increased dilution arising from runoff (1 ha) and groundwater recharge from the landfill mound. This dilution together with natural attenuation decreases the leachate levels in the aquifer.

4.5 HYDROLOGY

The hydrology of the area is dominated by Blue Angle Creek, Crooked River and Seven Mile Beach (Figure 2).

The closest environmental receptors of water running through the GWDD are Blue Angle Creek, Seven Mile Beach and Crooked River Estuary (Figure 1). Blue Angle Creek flows into the Crooked River Estuary at the northern end of Seven Mile Beach. The estuary discharges into the ocean when the mouth is open.

Previous Crooked River and Blue Angle Creek surface water quality investigations were discussed in the URS report (2002). The results indicate a considerable variation in water quality, particularly between dry and wet conditions, which may be associated with inputs from the wider catchment area.

When the entrance to Crooked River is open, the lower part of the river is well flushed with oceanic waters. This results in levels of nutrients, bacteria and toxicants that generally comply with guideline levels. Immediately following wet weather, water quality in the estuarine section of Crooked River generally deteriorates, with increased levels of particulate material, bacteria, sulphide, nutrients and metals (URS, 2003).

Of the four main tributaries that feed into the estuarine section of Crooked River, water quality in Blue Angle Creek was the most degraded with phosphorus, nitrogen, hydrogen sulphide, copper and zinc (URS, 2003).

All surface water runoff from the landfill mound is diffused into the surrounding sandy soils.



5 PREVIOUS MONITORING RESULTS

The primary conclusion from the monitoring report submitted by URS (2002-2003) prior to remediation in 2005 is summarised below:

 High concentrations of nutrients, in particular ammonia-nitrogen, continue to be detected under the site and migrating offsite. The levels recorded are well in excess of ANZECC (2000) guidelines for the protection of fresh and marine water ecosystems.

The following key points are also noted:

- High concentrations of ammonia-nitrogen and TKN were consistently detected in monitoring wells MW-1 and MW-5, which are located on the outer extent of the facility. This indicates the potential for migration of the nutrient plume in a south-easterly direction towards Seven Mile Beach. High concentrations of ammonia-nitrogen were also detected in monitoring wells MW-3 and MW-7, which are also located on the outer extent of the facility, indicating the potential for migration of the nutrient plume in a north-westerly direction towards Blue Angle Creek.
- Elevated concentrations of ammonia-nitrogen were detected in the shallow monitoring wells MW-9 to MW-11, adjacent to Blue Angle Creek.

6 SURFACE AND GROUNDWATER MONITORING

Surface and groundwater monitoring between May 2014 and February 2015 was undertaken by ALS Environmental. Sampling was carried out on the following dates:

- 27 May 2015
- 24 August 2015
- 4 November 2015
- 16 February 2016

The recommended procedure for sample collection, storage, handling and quality control generally employed by Ecowise (ALS) is outlined in the SGMP by AWT (1999a). E2W used the November 2008 sampling round to instruct Council staff on surface and groundwater sampling protocol. The samples were sent to ALS (Sydney) for laboratory analyses. E2W understand that ALS carry out the quarterly sampling at the GWDD and submit the samples to ALS (Sydney via the Wollongong office) for analyses.

6.1 MONITORING LOCATIONS

The following outlines the nature of the monitoring and analytical program at the site and the conditions at the time of sample collection from information provided by ALS and/or Council.

Groundwater was sampled from six monitoring wells (MW-3, MW-4, MW-5, MW-9, MW-10, MW-11), and three nested wells (i.e. MW-1S/MW-1D, MW-6S/MW-6D and MW-7S/MW-7D, where S = shallow, D = deep).



Surface water was sampled at two locations (ML-2 and ML-5) along Blue Angle Creek (Figure 2). Samples were not collected from ML-1, ML-3 or ML-4 due to restricted access to sample locations (private land owned by CB). Samples were not collected from Seven Mile Beach (BS-1 to BS-4), or the ocean (OS-1 to OS-4) during May 2012 to February 2013 (neither during 2014 to 2016). Previous results from these locations indicate that it is unlikely landfill leachate is impacting these areas.

Variable EC levels in surface water testing locations indicate that sampling may not have been undertaken during low tide (sampling at low tide provides a reflection of groundwater discharge).

6.1.1 Sampling Sites - Groundwater

The sampling sites are described below, while sampling depths for the bundled piezometers and conventional wells are summarised in Table 6.2.

- MW-1S (shallow 6 m depth) and MW-1D (deep 10.5 m depth)⁴ Located next to the previous multilevel piezometer MW-1 and approximately 40 m from the SE corner of the landfill perimeter. The well is situated down-gradient of the landfill mound and intended to intersect flow heading towards Seven Mile Beach.
- MW3 Approximately 20 m to the north of the landfill clearing, fronting native bushland. This well is to establish background water quality conditions and determine offsite migration of groundwater in a northerly direction.
- MW4 Located immediately adjacent to the night soil depot, which is a source of potential nutrient and bacterial contamination.
- MW5 Approximately 30 m to north of the night soil depot. The well is within the extent of contamination arising from the depot and landfill-impacted groundwater.
- MW-6S (shallow 6 m depth) and MW-6D (deep 10.5 m depth) Located next to multilevel piezometer MW-6 and approximately 50 m NW of landfill and night soil depot. The well is situated down-gradient of the landfill and night soil depot.
- MW-7S (shallow 6 m depth) and MW-7D (deep 10.5 m depth) Located 15 m to the east of multilevel piezometer MW-7 and approximately 100 m NW of landfill and night soil depot. This well was installed to establish background water quality conditions and determine offsite migration of groundwater contaminants towards Blue Angle Creek.
- MW9 Located offsite and adjacent to Blue Angle Creek downstream of ML-1 to determine if potentially contaminated groundwater is discharging into the estuarine environment.
 - MW10 Located offsite and adjacent to Blue Angle Creek downstream of ML-1 to determine if potentially contaminated groundwater is discharging into the estuarine environment.
 - MW11 Located offsite and adjacent to Blue Angle Creek downstream of ML-1 to determine if potentially contaminated groundwater is discharging into the estuarine environment.

⁴ Nested shallow and deep wells are constructed with a 3 m well screen.



6.1.2 Sampling Sites - Surface Water

Blue Angle Creek

- ML-1 Approximately 100 m upstream of the depot along Blue Angle Creek at the end of the tidal limit. This sampling location was chosen to establish upstream water quality and offsite conditions.
- ML-2 Approximately 500 m downstream of the depot along Blue Angle Creek. This sampling location was chosen to establish offsite and downstream water quality and assess the potential for contamination associated with the depot.
- ML-4 Approximately 100 m upstream of the flood gates along Blue Angle Creek. This sampling location was chosen to establish upstream water quality and offsite conditions.
- ML-5 Approximately 400 m downstream of the flood gates along Blue Angle Creek, between MW-9 and MW-11. This midstream sampling location was chosen to establish offsite receptor water quality conditions.

The surface water and groundwater sample locations are illustrated in Figure 2, and Table 6.1.2.

Table 6.1.2: Monitoring Summary for May 2015 to February 2016

Sample ID	Screen Interval (m AHD)	27 May 2015	24 August 2015	4 November 2015	16 February 2016
	& Sample Location	2015	2010	2012	2010
MW-1S	Approx. 0 to -3	X	Dry	X	Dry
MW-1D	Approx4 to -7	X	X	X	X
MW-3	0 to 1.5	X	X	X	X
MW-4	0.79 to -0.71	X	X	X	X
MW-5	0.55 to -0.95	X	X	X	X
MW-6S	Approx 0 to -3	X	Dry	Dry	Dry
MW-6D	Approx -4 to -7	X	X	X	X
MW-7S	Approx 0 to -3	X	X	X	X
MW-7D	Approx -4 to -7	X	X	X	X
MW-9	-0.53 to -1.53	X	X	X	X
MW-10	-0.525 to -1.525	X	X	X	X
MW-11	0.095 to -0.905	X	X	X	X
ML-1	Upstream of landfill	No access	No access	No access	No access
ML-2	Downstream of landfill	X	X	X	X
ML-3	Upstream of landfill	No access	No access	No access	No access
ML-4	Upstream of landfill	No access	No access	No access	No access
ML-5	Opposite landfill	X	X	X	X

Notes:

 $X = Sample\ collected.\ MW = Monitoring\ well\ sample\ from\ landfill\ site.\ ML = Surface\ water\ sample\ from\ Blue\ Angle\ Creek.\ The\ 6\ wells\ (MW-1S/MW-1D\ etc)\ are\ considered\ more\ reliable\ monitoring\ locations\ (compared\ to\ multilevel\ piezometers)\ as\ they\ were\ constructed\ with\ standalone\ 50\ mm\ diameter\ PVC\ screens\ and\ not\ the\ 7\ mm\ poly\ tubing\ (low\ flow\ system).$

e2W



6.2 SAMPLE COLLECTION AND LABORATORY ANALYSIS

The surface and groundwater analytical program from May 2015 to February 2016 is presented in Tables GW-1 and SW-1. The nested wells (MW-1S, MW-1D, MW-6S, MW-6D, MW-7S, MW-7D) installed in 2006 have replaced the bundled piezometers ⁵ (i.e. MW-1, MW-6, MW-7). The results from the nested wells are graphed separately for the water quality trend assessment.

7 MONITORING RESULTS

All groundwater and surface water analytical results from 2003 to 2016 are presented in Tables GW-1 and SW-1, with the most recent (4) monitoring data highlighted. The field records and laboratory reports are presented in Appendix A and Table 6 (summary of data set).

Compliances exceedances are noted in Appendix C (ML-5 and ML-2).

A summary of all available monitoring data (2003 to 2016) is presented in Graph-01 to Graph-07. The graphs illustrate ammonia and total phosphorous concentrations (key indicators of leachate impact) for the groundwater wells⁶ and surface water sampling locations as well as the depth to groundwater (m AHD, 2001 to February 2016).

The graphs illustrate the ammonia concentrations in the shallow (Graph 1) and deep monitoring wells separately (Graph 2). Contaminant migration rates and flushing characteristics are different at shallow and deep levels of the aquifer.

The graphs highlight water quality trends with respect to seasonal and water level changes, as well as water quality improvements associated with the landfill rehabilitation completed in 2005.

7.1 GROUNDWATER DATA

Groundwater was collected from a network of twelve monitoring wells at onsite and offsite locations (Figure 2) in May, August, November 2015 and February 2016. The results of the groundwater results obtained are summarised in Table GW-1, Graph-01 to Graph-05 and the following subsections.

7.1.1 Groundwater Depth and Flow Regime

The depth to groundwater was measured prior to each sampling event (in conventional wells) using a water level probe. The depth to the groundwater below top of casing and relative to a common reference (i.e. Australian Height Datum, m AHD) is presented in Table GW-1. The inferred groundwater contours are presented in Figures 3A & 3B. Reduced groundwater levels (m AHD) between 2001 and 2016 are illustrated in Graph-05.

⁵ The bundled wells were believed to provide spurious results due to the low purge volumes.

⁶ Results from multilevel piezometers MW-1, MW-6 and MW-7 (Graph-01 & Graph-03) are considered anomalous.



A groundwater divide is interpreted to occur at the landfill mound (sand dune area) and inferred to be located midway between Crooked River and Seven Mile Beach. As the position of the groundwater divide is influenced by the surface water bodies, the prevailing climate and recharge through the waste mound, it is likely local groundwater flow characteristics have altered over recent years.

The reduced groundwater levels from the twelve wells indicate a relatively moderate water table elevation. Field sampling records show that the depth to groundwater between October 2003 and February 2012 is below 1.87 mAHD. The groundwater levels recorded from the 2011-2012-2013, 2014-2015 monitoring round are similar to slighter lower than the 2015-2016 monitoring period, reflecting variable rainfall patterns and leachate movement over time. The 2015-2016 wet weather and higher water table is similar to the February 2011 wet period.

The inferred groundwater contours for the site are presented in Figures 3A & 3B (reflecting dry and wet weather periods with corresponding shallow and deep water tables). The groundwater levels and degree of mounding is generally variable, indicating a dynamic groundwater environment dependent upon rainfall recharge (aquifer is unconfined and sandy soils are highly permeable at the site).

7.1.2 Field Parameters

The groundwater, field parameters measured during sampling are considered indicative only (small purge volumes). Insitu measurements (within borehole) are likely to provide a more accurate rendition of the field chemistry, especially with respect to dissolved oxygen.

7.1.2.1 pH (field) and Redox

The groundwater pH measured from the twelve wells ranges from pH 5.1 to 8.1 (MW-10 & MW-7s). The pH in each well was generally stable over the four monitoring rounds (May 2015 to February 2016).

On its own, pH is not considered a reliable indicator of leachate contamination, as sediments and decomposing organic material associated with the creek bed also have a significant influence on pH.

Redox was not measured during the previous monitoring periods (2010 to 2011 and 2009- 2010) but was measured in August 2011 during 2011-2012 sampling round, and August 2012 during the 2012-2013 monitoring period. Similarly redox was only measured in August 2014 during the 2013-2014/2014-2015/2015-2016 sampling rounds, ranging from 130 to -138 mV (MW-7s and MW-9). In each well redox is generally comparable to previous monitoring periods.

7.1.2.2 Total Dissolved Solids (TDS and EC)

The TDS levels in groundwater collected from the site range from 162 to 6,840 mg/L (fresh to brackish). The lowest and highest TDS levels were obtained from MW-5 and MW-9, respectively. The TDS level for MW-9 is related to the well's proximity to Blue Angle Creek and associated tide and estuary mouth closure influences.



Salinity increases as groundwater flows towards Blue Angle Creek (MW-9 to MW-11). A decrease in salinity (TDS) occurs in several wells close the landfill perimeter (i.e. MW-3, MW-4, MW-5), which is interpreted to occur from dilution arising from an increase in stormwater runoff from the landfill mound post-capping (& decreased leachate generation).

7.1.2.3 Dissolved Oxygen (DO)

Field analyses from the twelve wells recorded dissolved oxygen (DO) concentrations ranging between 1.47 and 3.9 mg/L (MW-6D and MW-7s). The concentrations of DO reported for the site from 2003 to 2016 are variable. However, measurements may reflect the instruments (imprecise) used and/or purging process.

Based on the distribution of DO in the groundwater at the site, it is inferred that landfill leachate caused a depletion of groundwater DO and is generally increasing due to decreasing nutrient concentrations. This phenomenon is seen on many landfill sites, where organic carbon and nutrients provide surplus electron acceptors, which react with and consume the available DO in the groundwater. It is likely the groundwater under the landfill mound is anaerobic due to the presence of the landfill leachate (DOC, ammonia etc.) and poor flushing due to the landfill capping.

7.1.3 Nutrients

7.1.3.1 *Nitrogen*

Groundwater collected from the monitoring wells at the GWDD were analysed for ammonia-nitrogen, total Kjeldahl nitrogen (TKN) and oxidised nitrogen (nitrate and nitrite). Discussions regarding potential impact to the environment will focus on ammonia-nitrogen, as it is the main indicator of groundwater contamination from leachate.

The guidelines for total ammonia-nitrogen for the protection of fresh water and marine ecosystems vary according to pH and temperature. Given the range of pH and temperature measured across the site and in Blue Angle Creek, the guidelines are 1.88 and 2.84 mg/L for fresh and marine waters, respectively (at a pH of 7.3).

At least one groundwater samples collected from MW-3, MW-5, MW-6D, exceeded the ANZECC (2000) trigger value for ammonia. These monitoring wells are located adjacent or west of the landfill mound, indicating leachate is migrating towards Blue Angle Creek. The ammonia (20.5 mg/L in February 2016 & TKN=58.1 mg/L) reported in MW-03 is suspected to be an anomaly and should be re-sampled for quality control.

Groundwater wells (MW-1, MW-3 and MW-4 pre-rehabilitation) initially reported the highest concentrations of ammonia. Following the landfill rehabilitation, ammonia levels have declined in the shallow groundwater system and are below ANZECC guidelines (Graph-01, except MW-03 in February 2016 which is inferred as an anomaly). Groundwater from deep wells (MW-1D, MW-6D and MW-7D) show a clear declining trend, and now cycling above/below the ANZECC guidelines.



The ammonia concentrations show a clear reduction in nutrient loading in the deep aquifer (ammonia generally below 5 mg/L in the 3 key deep wells) and water quality improvement at the site since landfill closure and rehabilitation.

Nitrate was analysed for all samples (May 2015 to February 2016) with MW-1D, MW-6S, MW-6D, MW-7D, MW-9, MW-10 are generally below or marginally exceeded the ANZECC (2000) guideline (fresh water trigger value, 0.7 mg/L). The result for MW-1D (February 2016= 1.87 mg/L) is the highest result and slightly above the guideline.

All groundwater samples analysed from all twelve wells in May, August, November 2015 and February 2016 exceeded ANZECC (2000) fresh and marine water TKN trigger values (0.5 and 0.12 mg/L, respectively). Monitoring wells located adjacent to Blue Angle Creek (i.e. MW-9, MW-10, MW-11) continue to report stable or decreasing concentrations of ammonia (Graph 1). Table 7.1.3 (below) and Figure 4 provide an overview of groundwater ammonia trends from May 2015 to February 2016 reporting period.

Table 7.1.3: Groundwater Ammonia Trends - May 2015 to February 2016

Well ID	Ammonia Trend	Exceedance of ANZECC (2000) Ammonia	Trigger 20% exceedance (Ammonia)	Comment
MW-1S	Decreasing trend. below ANZECC	No exceedance	No exceedance	Shallow sample - east of landfill
MW-1D	Variable but decreasing trend below ANZECC (3.78 mg/L)	Exceedance	No exceedance	Deep sample - east of landfill
MW-3	Decreasing/stable trend. below ANZECC (anomalous February 2016 @20.9 mg/L)	Exceedance #	No exceedance	North of landfill. Re-sample of Feb 2016 required
MW-4	Decreasing/stabile trend. below ANZECC	No Exceedance	No exceedance	West of landfill
MW-5	Stable and below ANZECC	No Exceedance	No Exceedance	North of landfill.
MW-6S	Stable trend below ANZECC	No Exceedance	No Exceedance	Shallow sample - down- gradient of night soil deposit
MW-6D	Stable/decreasing trend. levels above and below ANZECC	Exceedance	No Exceedance	Deep sample - down-gradient of night soil deposit
MW-7S	Decreasing/Stabilising trend. below ANZECC	No exceedance	No exceedance	Shallow sample - down- gradient and adjacent to Crooked River Road
MW-7D	Decreasing/Stabilising trend. levels below and above ANZECC (6/5/2014= 1.91 mg/L)	Exceedance	No Exceedance	Deep sample - down-gradient and adjacent to Crooked River Road
MW-9	Stable/decreasing trend. below ANZECC	No exceedance	No exceedance	Next to Blue Angle Creek
MW-10	Stable/decreasing trend. below ANZECC	No exceedance	No exceedance	As above
MW-11	Stable/decreasing trend. below ANZECC	No exceedance	No exceedance	As above

Note: # inferred anomalous result. Three wells (MW-9, 10, 11) are located on the creek bank, potentially affected by flood waters and vegetation. ANZECC (2000) refers the marine trigger value (2.84 mg/L).



As outlined in the revocation notice, ammonia concentrations greater than 20% above ammonia concentrations reported in Table GW-1 of *Kiama Municipal Council, Gerroa Waste Disposal Depot - Annual Groundwater and Surface Water Monitoring Report - August 2006 to May 2007*, dated 17 August 2007 are to be highlighted (refer to Appendix C).

7.1.3.2 Total Phosphorus (TP)

The ANZECC (2000) TP guideline for fresh and marine ecosystems is 0.05 and 0.025 mg/L, respectively. Between May 2015 and February 2016, all groundwater sample results exceeded the ANZECC (2000) trigger values (Table GW-1 and Graphs 3 & 4).

MW-6D (located down gradient of the former night soil deposit and landfill) reported a maximum of 6.84 mg/L in the 2011-2012 monitoring period, however decreased to a maximum of 4.8 mg/L in the 2012-2013 monitoring period. In 2013-2014 and 2014 -2015 total phosphorous was a maximum of 9.07 mg/L and 3.93 mg/L, respectively indicating variability. In November 2015 the TP was reported at 12 mg/L at MW-6D, which is the highest to date (inferred as anomalous and requires re-sampling to verify). MW-7D reported a maximum (TP =8.14 mg/L in November 2015) which is similar to previous years (TP =8.46 mg/L, November 2014). TP at MW-6D and MW-7D (near former night soil) shows variability (Graph-4) and may relate to the 2013 dewatering/sludge pond decommissioning (further monitoring required to verify impacts in the deep groundwater) and high rainfall events during 2015 (April).

During the 2013-2014-2015-2016 monitoring period, MW-6S (shallow nested well) initially showed a decreasing trend and a maximum of 0.4 mg/L (May 2015), 0.8 mg/L (February 2014), comparable to the 2012-2013 maximum (0.63 mg/L) and significantly lower than the 2011-2012 maximum of 8.81 mg/L. TP at MW-4 (1.97 mg/L in November 2015, and located south of the night soil deposit) indicate an elevated, variable and potentially stabilising trend (Graph 3). The August 2012 & 2015 monitoring event reported a maximum TP (4.29 mg/L and 1.81 mg/L) at MW-4, which is lower than the maximum TP in 2011-2012 (7.98 mg/L, the highest to date since sampling began in June 2003).

The removal of the liner from the sludge pond in October 2011 coupled with high rainfall events may have impacted the subsequent results.

The concentrations of TP immediately east and north of the landfill (MW-3 and MW-5) and next to Blue Angle Creek (MW-9 to MW-11) are lower, indicating that the former night soil deposit is a likely source of TP.

In relation to the former night soil deposit (primary TP source), a localised TP plume is interpreted to potentially migrate towards Blue Angle Creek (MW-4/MW-6S to MW-7S, and to MW5 (TP= 1.39 mg/L in August 2015). The TP plume is also detected at MW-4 (south of the night soil deposit).

Potentially increasing TP concentrations in the deep wells (10.5 m depth, MW-6D, MW-7D) show that the plume may have reached the deep aquifer, however further monitoring is required to determine trends and some potentially anomalous results (e.g. TP=12 mg/L in November 2015). The TP concentrations of TP on the east/north (MW-3/MW-1) are likely to originate from the landfill mound.



TP concentrations at MW-1S (eastern side of landfill) show a variable trend; however further monitoring is required to verify the trend.

The TP results show variability and likely to be linked to flushing and desorption/sorption from the sandy aquifer matrix following from high rainfall events.

7.1.4 Hydrogeochemical Indicators

Concentrations of major ions (i.e. chloride, sulphate, calcium, magnesium, sodium, alkalinity and potassium) are presented in Table GW-1. The concentrations at all monitoring wells are within previously reported ranges and characterised by the ions sodium, chloride and bicarbonate (alkalinity).

The landfill is interpreted to contribute some concentrations of ions including calcium, potassium, magnesium and bicarbonate/alkalinity. Contribution of sodium and chloride is difficult to ascertain as these ions are common in the marine environments (e.g. salt spray, tidal influence) and abundant in wells close to Blue Angle Creek (MW-9, MW-10, MW-11) or in proximity to Seven Mile Beach (MW-1S, MW-1D).

7.1.5 Inorganic Contaminants (Iron, Manganese and Fluoride)

Iron concentrations were only analysed for samples collected in August 2015. Concentrations of iron (filtered at the laboratory) ranged between <0.05 and 1.28 mg/L (MW-11 is the maximum). Two wells (MW-7s, MW-11) reported concentrations were above the ANZECC (2000) guideline for iron in fresh water ecosystems (0.3 mg/L). Multiple natural and landfill related sources of iron are likely to exist at the site and offsite area (lithology and landfill).

The ANZECC (2000) guideline for iron is an indicative interim working level (IIWL) and is of low reliability. No guideline is available for iron in marine water, which is more relevant for Blue Angle Creek and Crooked River receiving water bodies.

Manganese concentrations were analysed for samples collected in August 2015. Concentrations of manganese ranged between 0.004 (MW-9) and 0.086 mg/L (MW-4), without exceeding the ANZECC (2000) fresh water guideline (1.9 mg/L).

The levels of filterable iron and manganese are generally similar to previous reporting periods. Variation in the concentrations may also reflect turbidity of water samples and filtering procedures.

Concentrations of fluoride (analysed in August 2015) ranged from <0.1 to 0.4mg/L (MW-6D), which are similar to previous reporting periods (2010-2014). No reliable ANZECC (2000) guideline exists for fluoride in fresh or marine waters.



7.1.6 Organic Contaminants

Dissolved organic carbon (DOC) concentrations were only analysed for samples collected on the August 2015. The concentration of dissolved organic carbon (DOC) in samples from the twelve wells ranged from 3 mg/L (MW-4) to 102 mg/L (MW-11). The results are generally comparable to previous monitoring periods, with lower concentrations noted in this period. No recommended ANZECC (2000) guidelines exist for DOC, but can be used to indicate organic carbon related to landfill leachate.

7.1.7 Discussion and Trends - Groundwater

The key trends in groundwater levels and nutrient contamination from 2003 to 2015 are presented in Graph-01 to Graph-05. The monitoring data indicates that ammonia concentrations in the deep groundwater are close to ANZECC guidelines (i.e. results at above and below ANZECC as shown in Graph-02). Trends for the deep wells (MW-1D, MW-6D and MW-7D) show a clear declining trend since August 2009 (Graph-02).

Groundwater (ammonia) trends for the three wells (MW-09, 10, 11) located adjacent Blue Angle Creek show a declining trend (Graph-01). These three wells are influenced from tides and flooding (including estuary mouth closures).

Graph-01 indicates ammonia concentrations in the shallow wells have steadily decreased (generally below guidelines) since land-filling operations at the GWDD ceased in October 2003.

Prior to landfill rehabilitation, groundwater quality trends indicate landfill leachate generation may be related to rainfall recharge into buried waste and subsequent groundwater and contaminant migration. The results post-landfill rehabilitation indicates landfill leachate concentrations in the shallow groundwater are decreasing, becoming diluted from attenuation/rainfall via runoff from the landfill mound. The potential for landfill leachate generation was significantly reduced following remedial works, as the buried waste was capped with an impervious clay barrier.

The groundwater ammonia trends (ammonia being a key landfill leachate indicator) indicate shallow groundwater quality is improving. The three deep wells installed in 2006 indicate the leachate plume in the deep parts of the aquifer is also improving (Graph-02) more slowly as flushing is lower at deeper levels in the aquifer (i.e. below sea level and across the groundwater divide).

7.2 SURFACE WATER

Surface water sampling was undertaken in May, August, November 2015 and February 2016. Samples were collected from two locations (ML-2 and ML-5, permission for accessing other locations was denied by site owner) locations along Blue Angle Creek (Figure 2).

Samples were not collected from ML-1, ML-3 or ML-4 due to restricted access (i.e. land is owned by Cleary Bros with access denied for sampling); therefore upstream water quality is relatively unknown and may be degraded due to farming and areas of acid soils. Sample locations, ML-1 ML-2 (downstream) and ML-5 (midstream) are not considered appropriate to assess water quality impacts



from the landfill and potential upstream sources. All analytical results for surface water monitoring points ML-2 and ML-5 are presented in Table SW-1.

Blue Angle Creek is tidally influenced and has a marine water influence at all sample locations, as shown by the broad range of TDS/conductivity results (i.e. fresh to saline, Table SW-1).

The surface water monitoring data at upstream/downstream locations is variable and likely to reflect a combination of tidal sampling regimes and inputs from the broader catchment area (e.g. samples should be coordinated with the tide so that both creek samples are collected during a run-out tide when the maximum amount of groundwater (potential leachate) discharges into the creek).

7.2.1 Field Parameters

7.2.1.1 pH (field) and Redox

The pH was similar at ML-2 and ML-5, and ranged from pH 6.5 to 7.3. Sampling results indicate that pH is slightly more acidic at upstream areas with Blue Angle Creek and may relate to the acid soils in the area.

Redox was only measured in August 2015 during 2015-2016 monitoring period, at ML-2 (-17 mV) and ML-5 (95 mV).

7.2.1.2 Total Dissolved Solids (TDS and EC)

Restricted access has limited the assessment of upstream and downstream locations. Previous monitoring period indicated that between August and November 2008, the concentrations of TDS at the Blue Angle Creek upstream location (ML-1) were less than those recorded downstream (ML-2). The TDS concentration upstream of the flood gates (ML-4) was less than those recorded downstream of the flood gates (ML-1 and ML-2).

Samples collected between May 2015 and February 2016 at the downstream location ML-2 were brackish to saline 7 (TDS = 2060 to 17,200 mg/L), while samples from the midstream location ML-5 were slightly fresher (TDS = 538 to 6,860 mg/L). The surface water samples are located in an area of the creek that is known to be influenced by tides. The presence of mangroves and other aquatic plants also reflects the typically saline water in the lower section of Blue Angle Creek.

Groundwater samples collected from MW-9, MW-10 and MW-11 were less saline than surface water samples collected from Blue Angle Creek and more saline than samples from all other groundwater monitoring wells sampled between May 2015 and February 2016. This data indicates that tidal waters from Crooked River Estuary can influence water quality and salinity of the 3 wells (MW-9, MW-10 and MW-11). Higher salinity reduces (as more marine water) the possibility of detecting leachate derived from the landfill.

.

⁷ Possibly reflecting collection of samples during high and low tide and stormwater runoff and rainfall



7.2.2 Nutrients

7.2.2.1 *Nitrogen*

Concentrations of ammonia in the surface waters collected from Blue Angle Creek have been, and continue to vary with time (Graph-06 and Table 7.2.2). The upstream catchment area of Blue Angle creek is improved pastures and grazing (i.e. agricultural sources of nutrients).

Table 7.2.2: Surface Water Ammonia Trends - May 2015 to February 2016

Sample	Minimum	Maximum	Trend	Comments &
ID	(mg/L)	(mg/L)		Trigger 10% Exceedance of Ammonia
ML-1	-	-	-	No site access
ML-2	0.08	1.76	Variable,	Yes. 10% Exceedance (1.38 mg/L @Nov
	(May 2015)	(Nov 2015)	generally	2015):
	-		decreasing	All below ANZECC guidelines
ML-3	-	-	-	No site access
ML-4	-	-	-	No site access
ML-5	0.83	2.39	Variable,	Yes. 10% Exceedance Aug 2015 (2.39 mg/L)
	(May 2015)	(Aug 2015)	generally	Below ANZECC guidelines, except for Aug
			decreasing	2015 (2.39 mg/L)

Note: ML-5 was sampled for the first time in four years in November 2008.

Highest ammonia (2.39 mg/L) concentration is associated with high TKN (2.6 mg/L), brackish water (TDS= 1370 mg/L) which is inferred to reflect background water quality (downstream of fertilised farms, sewerage treatment plant and former landfill).

Sampling at ML-5 (resumed in November 2008 but had not been sampled since October 2004) indicates that the water quality is variable (Table SW-1), indicating multiple pollutant sources and tidal influence.

Restricted access to upstream locations (ML-1, ML-3 and ML-4) limits conclusions regarding impacts due to the absence of upstream sample locations. Elevated ammonia in surface water compared to groundwater wells (& variable TKN concentrations) in proximity to the creek (MW-9, MW-10, and MW-11) indicates that sources of ammonia also occur from the upstream catchment area (agricultural land).

Concentrations of TKN exceeded the ANZECC (2000) guidelines for fresh and marine waters for all samples collected along Blue Angle Creek. The highest concentration was reported at the upstream location (ML-5, 3 mg/L in November 2015). Concentrations of TKN in groundwater are elevated and variable.

Increases in nitrogen from upstream and downstream of the landfill have been observed during previous monitoring periods. While these increases may be attributable to the discharge of ammoniarich groundwater from the landfill, other sources (random) of nitrogen input such as runoff from subcatchments and nutrients bound in sediments cannot be discounted. It is also possible that poor quality estuarine waters from Crooked River move up Blue Angle Creek during tidal cycles. Sewerage discharges into sand dunes may also occur due to capacity issues at the Gerroa sewerage treatment plant.



Interpretation of the nutrients into surface water bodies from the landfill is complicated by the sampling regime (i.e. sampling at various tides) and other potential sources of nitrogen. The fluctuating flow regime near Blue Angle Creek and wet weather events may reduce the potential for landfill leachate to impact the creek.

Total organic carbon (TOC) was not measured during the 2015 to 2016 monitoring period.

As stipulated in the revocation notice, ammonia concentrations greater than 10% above ammonia concentrations reported in Table SW-1 of Kiama Municipal Council, Gerroa Waste Disposal Depot -Annual Groundwater and Surface Water Monitoring Report - August 2006 to May 2007, dated 17 August 2007 are to be highlighted (Appendix C). Exceedances greater than 10% above ammonia were exceeded at ML-2 (trigger value = 1.38 mg/L@1.76 mg/L for November 2015), and ML-5 (trigger value = 2.38 mg/L@2.39 mg/L for August 2015). Based on the low ammonia results from monitoring wells (MW-9, MW-10, MW-11, Graph 1) in proximity to the creek, high ammonia concentrations in the surface water may be attributed to potential upstream sources (agriculture) or poor quality estuarine water (tidal or mouth closure).

7.2.2.2 Total Phosphorous (TP)

Concentrations of TP from Blue Angle Creek were analysed from all samples collected in May, August, November 2015, and February 2016 (Table SW-1).

Previous levels (2011-2012 monitoring period) reported an increase in TP concentrations, which exceeded the IIWL⁸ ANZECC 2000, (fresh 0.05 mg/L, marine 0.025 mg/L) at ML-2 and ML-5 for all sampling rounds. The 2012-2013 monitoring period reported an variable increase in TP concentrations which exceeded the IIWL⁸ ANZECC 2000, (fresh 0.05 mg/L, marine 0.025 mg/L) at ML-2 (0.05 mg/L, May 2012 and 0.13 mg/L, February 2013) and ML-5 (1.95 mg/L, February 2013). The 2013-2014 monitoring period reported two exceedences of the ANZECC 2000 guidelines for ML-2 in May 2013 & February 2014 (0.09 mg/L and 0.12 mg/L respectively) and one exceedence in ML-5 in 2013 (0.06 mg/L).

TP concentrations at ML-2 ranged from 0.03 mg/L to 0.16 mg/L and continue to show a variable trend for 2015-2016. The trend may be associated with the dynamic nature of the surface water and surrounding environment.

ML-5 reported all TP concentrations ranged from 0.03 mg/L to 0.11 mg/L in the 2015-2016 reporting period, indicating variability over time (Graph-07). The variability is considered to reflect the dynamic nature of the environment.

7.2.3 Bacteriological Contaminants

Surface water sample locations (ML-2 and ML-5) were analysed for thermotolerant (faecal) coliforms and enterococcus coliforms on 24 August 2015 (Table SW-1).

⁸ ANZECC (2000) Indicative Interim Working Levels (IIWLs).

⁹ Laboratory Level of Reporting (LOR)



Both samples results in August 2015 for enterococcus were above ANZECC (2000) fresh and marine guidelines (35 CFU/100 mL), downstream sample ML-2 reported 590 CFU/100 mL and midstream sample ML-5 reported ~240 CFU/100 mL. Both locations are significantly lower than 2010-2011 monitoring period results (ML-2, 1300 CFU/100 mL and ML-5, 1200 CFU/100 mL, November 2010) but above the 2011-2012 (ML-2, 18 CFU/100 mL and ML-5, 8 CFU/100 mL) and 2012-2013 (ML-2, 4 CFU/100 mL and ML-5, 15 CFU/100 mL).

Surface water samples reported levels of thermotolerant (faecal) coliforms below ARMCANZ (2000) guidelines (150 CFU/100 mL) for marine and fresh water ecosystems at ML-2 (640 CFU/100 mL) and ML-5 (\sim 1100 CFU/100 mL).

Multiple sources of coliforms exist in the surface water system, with the capped landfill representing an insignificant contribution, due to it's distance from the creek (i.e. local fauna and flora, farms) and filtering in the aquifer.

7.2.4 Inorganic Contaminants

Dissolved organic carbon (DOC) concentrations were not analysed during the 2012-2016 monitoring periods.

Surface water samples from August 2015 reported elevated concentrations of iron (ML-5 = 2.09 mg/L, ML-2 = 1.64 mg/L) above ANZECC 2000 guidelines (0.3 mg/L). However, the ANZECC (2000) guideline for iron in fresh water is a low reliability IIWL.

Manganese concentrations for ML-2 (0.023 mg/L) & ML-5 (0.032 mg/L) were reported below the ANZECC (2000) fresh water guidelines (low reliability IIWL).

7.2.5 Major Ions

Concentrations of major cations (sodium, potassium, alkalinity, magnesium, calcium) in the surface water (Blue Angle Creek) indicate domination of sodium (marine water influence), which is consistent with previous monitoring rounds.

7.2.6 Quality Assurance/Quality Control

Interpretive Quality Control Reports (QCI, Appendix A) provided by ALS (Sydney) of the surface and groundwater laboratory data were reviewed for the four sampling rounds (May, August, November 2015 and February 2016). No laboratory outliers or exceedances of holding times were noted during the 2015-2016 monitoring period.



8 LEACHATE PLUME AND LANDFILL REHABILITATION

The monitoring results have been used to assess potential impacts to fresh and marine aquatic ecosystems. The groundwater migrating from the former landfill discharges to Blue Angle Creek and Seven Mile Beach. The range of groundwater contaminants identified from the latest monitoring events indicates the GWDD is a source of leachate (mostly ammonia and TKN), total phosphorous and iron

Ammonia is the primary landfill leachate indicator. However, the waste is also a source of dissolved salts, metals and organics associated with the dissolution of ions (predominantly calcium and bicarbonate).

Following the closure of the landfill in 2003 and remedial works completed by Council and E2W in February 2005, the generation and migration of ammonia has declined in the shallow and deep groundwater system (Graphs-01 & 02). Monitoring wells MW-3, MW-4 and MW-5 are considered to reflect the typical groundwater quality arising from the landfill rehabilitation (>80% decrease in ammonia over time).

The deep groundwater monitoring wells (MW-1D, MW-6D and MW-7D) installed in 2006 show water quality improvements but at a slower rate relative to the shallow groundwater. Deep groundwater takes longer to improve due to the slower groundwater flushing. However, since August 2009 the ammonia concentrations in deep wells show a clear decreasing trend (Graph-02).

The most significant contaminant is ammonia-nitrogen, with a remnant plume extending in both north-west and south-east directions reflecting flows either side of the groundwater divide. Prior to rehabilitation, leachate originating from the landfill and night soil depot infiltrated the shallow aquifer, as well as migrating under the predominant groundwater flow regime towards Seven Mile Beach and Blue Angle Creek, respectively.

E2W consider the leachate plume in the shallow/deep groundwater has shrunk due to a decrease in leachate generation (capping of landfill causing an increase in runoff) as well as from natural attenuation processes (including dilution). The time series trends show that significant groundwater quality improvements occured after approximately 5 years.

Monitoring results indicate a clear improvement in water quality and aquifer restoration. Landfill rehabilitation is considered to have achieved a practical and successful outcome over a 10 year period.

Previous results (2008-2009, 2009-2010, 2010-2011, 2011-2012, 2012-2013, 2013-2014, 2014-2015) results from the three groundwater wells located along Blue Angle Creek (MW-9, MW-10 and MW-11) indicate ammonia concentrations below the ANZECC (2000) guidelines (with the exception of MW-11 in 2009-2010, Table GW-1). Recent 2013-2014/2014-2015/2015-2016 results continue to report a decrease in ammonia, which are below ANZECC (2000) guidelines.

Some elevated phosphorous occurs around the former night soil and is interpreted to arise from heavy wet weather and land disturbance around the source area (i.e. removal of lined ponds in ~2011).

The landfill rehabilitation conducted between July 2004 and February 2005 has resulted in a measurable improvement in the shallow groundwater quality at GWDD. While ammonia remains



elevated but has began to fall below the ANZECC 2000 guidelines in the deep groundwater, E2W consider that water quality will continue to improve in shallow/deep aquifer due to reduced leachate generation and ongoing natural attenuation.

The surface water results from the 2015-2016 monitoring period reported one guideline exceedences for ML-5 in November 2015 (2.39 mg/L). The surface water environment is dynamic and influenced by tidal flushing and discharges (runoff, seepage, baseflow) from the surrounding catchment and aquifer.

8.1 ECOLOGICAL ISSUES

Groundwater migrating from the landfill to Seven Mile Beach is diluted by the dynamic processes operating in this environment. Contaminants contained within this discharge may be diluted and dispersed via biological, chemical and physical processes occurring at the groundwater/salt water interface. Plant uptake of excess nutrients in the groundwater may also occur as the depth to groundwater becomes shallower as it approaches the beach.

The effect of nutrient-impacted groundwater discharging to Blue Angle Creek and/or Crooked River Estuary is unclear and difficult to ascertain given the variability, dynamic environment, and multiple nutrient sources in the catchment area.

Potential impacts of landfill leachate to Blue Angle Creek would depend on the groundwater-surface water interaction, climate and the rehabilitation works. Results from MW-9, MW-10 and MW-11 (monitoring wells adjacent to the creek) indicate a reduction in ammonia (Graph-01). Previously, MW-11 had a history of variable ammonia concentrations; however since May 2010 ammonia has remained below ANZECC guidelines.

E2W interpreted that some nitrogen-impacted groundwater would discharge to the creek (and consequently to the estuary), however the extent of attenuation of the nitrogen plume prior to discharge is unclear. Attenuation is likely to occur through a combination of dilution, mixing of groundwater from the creek, flows and tidal movements within the creek and estuary and oxidation of the ammonia to nitrate/nitrite and generation of nitrogen gas.

9 **CONCLUSIONS**

Surface and groundwater monitoring was completed at the GWDD by ALS in May, August, November 2015 and February 2016 for the EPL (2015). The data has been assessed by E2W to identify potential impacts to the groundwater and surface water systems. The following conclusions are offered:

• The rehabilitation of the landfill mound and night soil depot (completed February 2005) has demonstrated a measurable improvement of the local groundwater quality. The improvement to local surface water quality is not clear, and impacts are not readily discernible from landfill or background sources (agricultural, tidal water quality etc).



- Groundwater at the landfill site is directed towards Blue Angle Creek (base-flow discharge) and Seven Mile Beach (via a groundwater salt water interface).
- The key landfill indicator (ammonia-nitrogen) shows a decreasing/stabilising trend in shallow and deep wells located next to the landfill mound and former night soil deposit. Some data anomalies exist for the 2015-2016 monitoring period (e.g. MW-03 ammonia= 20.9 mg/L and TKN=58.1 mg/L in February 2016, and MW-6D with TP=12 mg/L in November 2015, and are to be addressed with future monitoring rounds).
- Elevated concentrations (above ANZECC 2000) of nutrients, in particular ammonia, continue to be detected in the shallow (MW-5) and deep groundwater (MW-6D, MW-7D with results cycling slightly above and below the ANZECC (2000) guidelines.
- Catchment area characteristics, climate and tidal regime all influence water quality in Blue Angle Creek and the adjacent wells (MW-9, MW-10 and MW-11).
- Concentrations of total phosphorous (TP) in the shallow/deep groundwater is generally variable, with elevated and variable trends at MW-3, MW-4, MW-5, MW-6D, MW-7D. Areas of variable TP is associated to the former night soil deposit, and likely mobilisation during heavy rainfall (e.g. January and April 2015).
- Ammonia within surface water samples collected at downstream locations (Blue Angle Creek) are generally variable and similar to previous years. An exceedance (greater than 10% above ammonia values in Appendix C) was noted at ML-2 for November 2015 (1.76 mg/L) and at ML-5 for August 2015 (2.39 mg/L). Based on the low (below guidelines) ammonia results from monitoring wells (MW-9, MW-10, MW-11, Graph 1) in proximity to the creek, ammonia in surface water may be attributed to potential upstream sources (agriculture) or poor quality estuarine water (tidal- and estuary mouth closures).
- Total phosphorus (TP) concentrations at ML-5 ranged from 0.01 to 0.12 mg/L in 2011-2012, however during 2012-2013, TP slightly increased and ranged from <0.01 to 1.95 mg/L. TP in the 2013-2014 monitoring round is comparable to that reported in 2011-2012 (0.01 mg/L to 0.06 mg/L), whilst TP ranges from 0.05 to 0.22 mg/L in 2014-2015. TP ranges from 0.03 to 0.16 mg/L in 2015-2016. TP in surface water is generally variable due to the dynamic nature of the tidal creek. It is likely that water quality in Blue Angle Creek reflects other nutrient sources in the catchment as well as from the GWDD.
- All other water quality indicators were consistent with the previous monitoring results.
- Monitoring results indicate a clear improvement in water quality and aquifer restoration at Gerroa.
 Landfill rehabilitation is considered to have achieved a practical and successful outcome over a 10 year monitoring period.

The nutrient concentrations (particularly ammonia) in the shallow and deep groundwater are likely to continue to decrease over time (note: phosphorous may take longer to decline as it is likely to adsorb/retarded by the aquifer matrix and mobilised by heavy rainfall). The landfill capping system reduces rainfall infiltration into the buried waste (reduces leachate generation) and diverts runoff from the 3 Ha capped mound into the aquifer, causing dilution and attenuation of the residual leachate.

Monitoring of surface and groundwater conditions at the GWDD following the completion of landfill remediation has provided beneficial data regarding the effectiveness of the rehabilitation works. The "surface and shallow" groundwater quality is showing signs of more frequently meeting the ANZECC (2000) guidelines. More consistent water quality and meeting of the guidelines is expected in the near future.



Based on recent groundwater trends, it is interpreted that the deep groundwater quality is beginning to meet (on some sampling events) the ANZECC (2000) guidelines for ammonia (key landfill leachate indicator) and demonstrating the effectiveness of landfill rehabilitation.

9.1 **RECOMMENDATIONS**

In order to improve the quality of monitoring at the site, E2W recommends that the following be incorporated into subsequent sampling rounds:

- Obtain survey details for the 6 piezometer wells (MW-1S/D, MW-6S/D and MW-7S/D).
- Assess quality assurance and control from laboratory and perform re-analyses for unusually elevated results.
- Perform reactive phosphorus (total & reactive) analyses at key wells (e.g. MW-6D, MW-7D).
- Review tidal charts and climate prior to surface water sampling. Sample collection should be coordinated with the tide so that both creek samples are collected during a low run-out tide when the
 maximum amount groundwater discharges into the creek.
- Further investigation of alternate sample locations to allow assessment of upstream water quality and the downgradient impacts associated with the landfill (i.e. replacement of previous locations @ ML-1, ML-3 and ML-4. It is noted that alternate locations are within private land "Cleary Bros site" and access has been denied).
- The proposed groundwater and surface water monitoring program for the GWDD is summarised in Table 3.1. The sampling methods are presented in Tables 1 and 2 of the Revocation of Licence (DECC, 29 May 2008).
- Discontinue bacteriological monitoring of surface water samples (i.e. not an EPL requirement and unlikely to indicate leachate discharges).



10 LIMITATIONS

Earth2Water Pty Ltd has prepared this report for the use of Kiama Municipal Council in accordance with the standard terms and conditions of the consulting profession. This report is prepared in accordance with the scope of work and for the purpose outlined in the proposal. The methodology adopted and sources of information used by E2W are outlined in this report.

E2W has made no independent verification of this information beyond the agreed scope of works and E2W assumes no responsibility for any inaccuracies or omissions.

This report was prepared in March 2016 and is based on the information reviewed at the time of preparation. This report should be read in full. No responsibility is accepted for use of any part of this report in any other context or for any other purpose or by third parties.

The precision with which conditions are indicated depends largely on the frequency and method of sampling and the uniformity of conditions as constrained by the project budget limitations. The behaviour of groundwater and some aspects of contaminants in soil and groundwater are complex. Our conclusions are based upon the analytical data presented in this report and our experience.

Where conditions encountered at the site are subsequently found to differ significantly from those anticipated in this report, E2W should be notified of any such findings and be provided with an opportunity to review the recommendations of this report.



11 REFERENCES

Australian and New Zealand Environment and Conservation Council & Agriculture and Resource Management Council of Australia and New Zealand (2000). *Australian and New Zealand Guidelines for Fresh and Marine Water Quality*.

Australian Water Technologies (2000). Surface Water and Groundwater Monitoring Program, Gerroa Waste Disposal Facility.

DUAP (1996). EIS Practise Guideline: Landfilling

Hazelton (1992). Soil Landscapes of the Kiama 1:1,000,000 Sheet, Department of Conservation and Land Management.

Landfill Management Services Pty Ltd (2003). *Landfill Gas Evaluation Gerroa Landfill*. *NSW EPA (1996) Environmental Guidelines: Solid Waste Landfills*.

DECC NSW (2007) *Guidelines for the Assessment and Management of Groundwater Contamination*, New South Wales Department of Environment and Climate Change, Sydney.

R.M. Koerner and T-Y.Soong, (1998). Analysis and Design of Veneer Cover Soils. Proceedings of 6th International Conference on Geosynthetics, Vol., pp.1-23, Atlanta, Georgia, USA.

URS Australia Pty Ltd (2002). Annual Surface and Groundwater Monitoring Report for the Gerroa Waste Disposal Facility, July 2001 to June 2002.

URS Australia Pty Ltd (2002). Annual Surface and Groundwater Monitoring Report for the Gerroa Waste Disposal Facility, July 2002 to June 2003.



Tables

Gerroa Waste Dispo	sal Depo	ot																										
Sample ID	ANZEC		MW1S	MW 1S	MW 1S	MW1S	MW1S	MW 1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1S	MW1D	MW 1D
Field Measurements	Fresh	Marine	21/5/10	17/8/10	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16	21/5/10	17/8/10
Ground Level (m AHD)			NA												-	-	-										NA	
Depth to Groundwater (m AHD)															-	-		_										
Groundwater depth (m bTOC)			3.74	3.52	2.88	3.3	3.25	3.17	3.53	3.86	3.33	3.6	3.9	3.75	3.25	3.16	3.74	Dry - no sample	3.29				3.06		3.2		3.62	3.39
Height of Stick up (m)			0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	Sample	0.65				0.65		0.65		0.62	0.62
Groundwater Depth (mbgl)			3.09	2.87	2.23	2.65	2.6	2.52	2.88	3.21	2.68	2.95	3.25	3.1	2.6	2.51	3.09		2.64				2.41		2.55		3	2.77
pH (field)	6.5-8.0 (a)	8-8.4 (a)	6.8	6.9	6.7	6.5	6.80	6.70	7.10	7.00	6.60	6.9	7.4	7	6.40	6.5	6.8						6.2		6		6.9	7.2
Temperature (T deg C)																-												
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		1.95	1.41	0.602	0.49	0.51	0.44	0.47	0.40	0.50	<1	0.464	1.1	0.41	0.42	0.548						772		342		2.4	2.43
Salinity (ppt)					-																							-
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	1.39	2.41	1.58	1.8	1.62	2.49	2.02	2.21	1.62	0.8	1.73	1.48	1.57	2.51	0.72	-	2.92				2.19		2.2		2.56	2.59
Dissolved Oxygen (%)					17.10							8.2																-
Turbidity (NTU)	6-50 (a)	0.5-10 (a)														-												-
Redox Potential (mV)			-	-	-	-	-	-52.2	-	-	-	< 0.1			-	-57	-										-	-
Comments			nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc										nc	nc
Sodium			-	-	54	-	-	21	-	-	-	22			-	42	-										-	-
Potassium					4			2				3				2												
Calcium				-	41		-	39	-		-	52			-	24											-	-
Magnesium				-	19	-	-	10	-	-	-	9			-	8	-											-
Chloride				-	113		-	38	-		-	34			-	51											-	-
Alkalinity (as CaCO3)			386	291	142	176	150	126	165	147	164	159		362	100	128	163		164				102		51		611	494
Bicarbonate			386	291	142	176	150	126	165	147	164	159		362	100	128	163	-	164				102		51		611	494
Carbonate (as CaCO3)			<1	<1	<1	<1	<1	<1	<1	<1	<1	<1		<1	<1	<1	<1		<1				<1		<1		<1	<1
Sulphate (SO4)				-	8	-	-	<1	-	-	-	6			-	3	-										-	-
pH (lab)				-	-		-	-	-		-				-	-			6.4									-
Total Dissolved Solids (TDS)			1300	982	430	416	436	386	350	250	390	346		640	362	309	442		516				734		385		1340	1330
Hardness (as CaCO3)				-	-		-	-	-		-				-	-												-
Total Suspended Solids (TSS)				-	-	-	-	-	-	-	-				-	-	-										-	-
Iron (filtered)	0.3 (1)		-	-	13.9	-	-	33.2	-	-	-	26.2			-	29.0	-											-
Manganese	1.90				0.046			0.05	-			0.051				0.050											-	
Nitrate (NO3 as N)	0.7 (7)		0.26	0.04	0.02	0.21	0.02	0.02	< 0.01	< 0.01	< 0.01	0.12		< 0.10	< 0.01	< 0.01	< 0.01		< 0.01				< 0.01		< 0.01		1.36	4.91
Nitrite (NO2 as N)			< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01		< 0.10	< 0.01	< 0.01	< 0.01	1	< 0.01				< 0.01		< 0.01		0.06	0.09
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	0.27	< 0.01	0.05	0.05	0.04	0.06	0.02	< 0.01	0.05	< 0.01		0.13	0.34	0.28	0.21	1	0.31				0.09		0.23		22.6	32.4
Total Kjeldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	4.4	46.8	2.6	1.9	2.50	0.90	1.10	2.20	1.2	1.0	Ì	1.8	1.9	1.3	4.6	-	2.7	1			3.5		2.1	Ť –	45.2	53.2
Dissolved Organic Carbon	(-/	1.77			38		-	31	-		-	26	i e			58		i										
Fluoride (Electrode)	1	1		-	0.3		-	0.2	-		-	0.3			-	0.2		i										
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.61	49.8	0.32	0.88	0.31	0.38	0.5	0.66	0.33	0.52		0.72	0.38	0.31	0.86	ĺ	0.44	1			0.26		0.18	Ť –	0.12	8.87
Note:	2.20(1)		01		02	00		00																				

Note: Exceeds ANZECC (2000) guidelines marine/fresh water ecosystems

Focus of this monitoring report nc = no comment NA = nc

35

- Notes:

 1. Trigger value is an indicative interim working level only (IIWL).

 2. Ammonia trigger at pH = 8.0, for a 95% protection, corrected for average pH = 7.3.

 3. Trigger value for oxides of Nitrogen (NOx) for towland rivers in NSW.

 4. Trigger value for oxides of Nitrogen (NOx) for smalrine ecosystems in NSW.

 5. Trigger value for total Nitrogen in lowland rivers in NSW.

- Trigger value for total Nitrogen in marine ecosystems in NSW.
 Trigger value for a 95% protection level.
 Guidelinefor water quality and aesthetics: primary contact.
 Reference only, not dirtectly applicable to groundwater.

Sample ID	ANZEC	C, 2000	MW 1D	MW1D	MW1D	MW 1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D	MW1D
ield Measurements	Fresh	Marine	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/10
Ground Level (m AHD)																-								
Depth to Groundwater (m AHD)																								
Groundwater depth (m bTOC)			2.77	3.16	3.12	3.04	3.5	3.73	3.2	3.48	3.77	3.6	3.1	3	3.64	3.54	3.15	3.65	3.68	3.54	2.92	3.84	3.09	3.6
Height of Stick up (m)			0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62	0.62
Groundwater Depth (mbgl)			2.15	2.54	2.5	2.42	2.88	3.11	2.58	2.86	3.15	2.98	2.48	2.38	3.02	2.92	2.53	3.03	3.06	2.92	2.3	3.22	2.47	2.98
oH (field)	6.5-8.0 (a)	8-8.4 (a)	6.8	7	7.30	7.60	7.60	7.50	7.80	7.5	7.5	7.5	7.30	7.5	7.5	7.5		7.6			7.5	7.7	7.3	7.3
Temperature (T deg C)																								
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		2.22	1.95	1.72	11.00	1.26	0.99	0.59	<1	0.657	0.695	0.63	0.624	0.624	0.59		0.467	0.622	0.636	707	709	803	631
Salinity (ppt)			-	-	-	-	-									-								
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	1.67	2.31	1.54	2.51	1.72	2.60	5.30	2.2	1.98	2.6	1.97	1.62	2.50	1.82	1.88	3.50	2.90	2.50	3.45	3.05	1.71	3.7
Dissolved Oxygen (%)			18.20		-	-	-			22.2								36.2						
Furbidity (NTU)	6-50 (a)	0.5-10 (a)	-	-	-	-	-									-								
Redox Potential (mV)			-	-	-	-91.6	-			< 0.1				-78		-		60				-37		
Comments			nc	nc	nc	nc	nc	nc	nc	nc	nc	nc												
Sodium			239	-	-	118	-			22				22				14				37		
Potassium			44	-	-	15				11				15				10				13		
Calcium			115	-	-	68	-			64				62				65				72		
//agnesium			64	-	-	27	-			23				24		-		20				24		
Chloride			488	-	-	146	-			12				20				21				37		
Alkalinity (as CaCO3)			231	345	297	316	322	363	279	288	293	292	284	316	298	254	283	225	280	296	272	262	291	252
Bicarbonate			231	345	297	316	322	363	259	288	293	292	284	316	298	254	283	225	280	296	272	262	291	252
Carbonate (as CaCO3)			<1	<1	<1	<1	<1	<1	21.00	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Sulphate (SO4)			160			34.00				6				6								10		
oH (lab)			-	-	-	-	-									-	7.20	7.60	6.40	7.20				
Total Dissolved Solids (TDS)			1420	1160	940	700	772	580	340	264	418	320	372	328	328	309	272	334	397	355	379	443	480	419
Hardness (as CaCO3)			-	-	-	-	-									-								
Total Suspended Solids (TSS)																								
ron (filtered)	0.3 (1)		9.93	-	-	0.10	-			< 0.05				< 0.05		-		< 0.05				< 0.05		
Manganese	1.90		0.036	-	-	0.008	-			0.001				0.010				0.00				0.006		
Nitrate (NO3 as N)	0.7 (7)		0.86	0.76	1.55	2.39	1.79	< 0.01	1.57	2	2.13	2.38	1.06	0.97	0.03	0.11	0.17	2.1	1.4	0.35	1.87	0.02	0.13	0.02
Nitrite (NO2 as N)	- \ /		0.12	0.02	0.08	0.01	0.06	< 0.01	0.01	< 0.01	< 0.01	< 0.01	0.08	0.07	< 0.01	< 0.01	0.03	<0.01	< 0.01	< 0.01	<0.01	< 0.01	< 0.01	0.01
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	25.7	6.67	2.56	0.12	2.15	3.16	0.07	0.09	0.17	0.26	0.27	2.88	4.13	2.9	1.29	0.02	1.3	1.51	< 0.01	0.28	3.78	2.43
Total Kieldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	31.3	9.6	4.60	1.70	3,30	4.00	1.50	1.4	1.9	1.2	1.10	3.4	4.4	3.1	1.6	0.6	2.1	1.8	0.6	0.7	4.6	3.3
Dissolved Organic Carbon	2.2 (0)	22 (0)	22			12	-		50	6				15				8			-10	9		0.0
luoride (Electrode)			<0.1			0.2				0.1			-	0.2				0.10		l		0.20		
	0.05 (7)	0.025 (7)	<0.01	0.35	0.03	0.15	0.04	0.26	0.08	0.18	0.16	0.02	0.03	0.02	0.44	0.1	0.0	0.10	0.3	0.1	0.05	0.46	0.06	0.17
otal Phosphorus (TP)																								

Gerroa Waste Dispo	sal Dep	ot																																
Sample ID	ANZEC		MW3	MW 3	MW 3	MW3	MW3	MW 3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW3	MW4	MW 4	MW 4	MW4	MW4	MW 4	MW4	MW4
Field Measurements	Fresh	Marine	21/5/10	17/8/10	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16	21/5/10	17/8/10	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11	1/2/12
Ground Level (m AHD)			4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.00	4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.79
Depth to Groundwater (m AHD)			0.37	0.59	1.18	0.83	0.87	0.95	0.48	0.25	0.79	0.49	0.25	0.40	0.88	0.97	0.35	0.12	0.83	0.35	0.30	0.42	1.06	0.29	0.90	0.42	0.88	1.08	1.69	0.99	1.31	1.41	0.98	0.76
Groundwater depth (m bTOC)			4.08	3.86	3.27	3.62	3.58	3.5	3.97	4.2	3.66	3.96	4.2	4.05	3.57	3.48	4.1	4.33	3.62	4.1	4.15	4.03	3.39	4.16	3.55	4.03	4.36	4.16	3.55	4.25	3.93	3.83	4.26	4.48
Height of Stick up (m)			0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45
Groundwater Depth (mbgl)			3.63	3.41	2.82	3.17	3.13	3.05	3.52	3.75	3.21	3.51	3.75	3.6	3.12	3.03	3.65	3.88	3.17	3.65	3.70	3.58	2.94	3.71	3.10	3.58	3.91	3.71	3.1	3.80	3.48	3.38	3.81	4.03
pH (field)	6.5-8.0 (a)	8-8.4 (a)	7	7.1	7	7.2	7.50	7.50	7.80	7.40	7.50	7.3	7.5	7.4	7.20	7.6	7.4	7.4		7.1			7.1	7.3	7.2	7	6.9	6.8	6.4	6.4	6.80	6.80	6.90	6.90
Temperature (T deg C)	0.0 0.0 (0)	0 011 (0)			-	-	-	-								-	-											-					-	
Electrical Conductivity (mS/cm)	0.125-2.2 (a)	,	0.74	0.74	0.754	0.48	0.56	0.70	0.47	0.46	0.39	<1	0.633	0.631	0.51	0.704	0.760	0.787		0.946	0.932	0.812	598	962	843	822	0.75	0.81	0.468	0.47	0.73	0.68	0.72	0.70
Salinity (ppt)			-		-	-	-	-	-	-			0.000		-	-	-			0.00.0	0.002	0.0.			0.0			-	-	-			-	-
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	3	1.91	2.67	1.68	4.50	2.07	1.58	1.76	2.66	0.7	1.75	1.94	1.68	1.84	1.54	2.36	2.69	1.4	2	3.8	3.05	3.21	1.54	2.8	2.55	2.32	1.46	2.21	1.78	3.09	2.64	2.34
Dissolved Oxygen (%)		, , ,			28.20	-	-	-				7				-	-			14.5								-	15.90	-			-	-
Turbidity (NTU)	6-50 (a)	0.5-10 (a)				-	-								-														-	-	-			-
Redox Potential (mV)					-	-	-	-81.6	-			< 0.1			-	-137	-			-114				-138			-	-	-	-	-	-41.5	-	-
Comments			nc	nc	nc	nc	-	-	-	-	nc	nc	nc	nc	nc	nc	nc	nc									nc	nc	nc	nc	-	-	-	-
Sodium					71			37				27				49				86				73					14.00			12		-
Potassium					3	-		2				3				3				3				3				-	1.00			2		-
Calcium					90			88				102				79				117				98				-	91.00			109		-
Magnesium					6	-	-	8	-			9			-	8	-			12				13			-	-	7.00	-	-	7	-	-
Chloride					127	-	-	53	-	-	-	130			-	92	-			205				106			-	-	23.00	-	-	25	-	-
Alkalinity (as CaCO3)			329	274	215	171	125	259	193	235	165	183	295	270	238	227	216	194	292.00	211	245	236	204	250	169	230	274	303	234	151	248	282	251	296
Bicarbonate			329	274	215	171	125	259	193	235	157	183	295	270	238	227	216	194	292.00	211	245	236	204	250	169	230	274	303	234	151	248	282	251	296
Carbonate (as CaCO3)			<1	<1	<1	<1	<1	<1	<1	<1	8.00	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Sulphate (SO4)					28			3				<1				6				11.00				13					11.00			10.00		-
pH (lab)						-													7.2	7.1	6.2	6.8						-	-					-
Total Dissolved Solids (TDS)			384	436	484	318	390	414	322	340	238	496	432	268	298	363	494	443	387	729	585	485	366	565	488	476	354	352	330	252	318	400	388	382
Hardness (as CaCO3)					-		-									-	-										-	-		-	-		-	-
Total Suspended Solids (TSS)										-	-																-	-		-	-			-
Iron (filtered)	0.3(1)				14.3			7.55		-	-	9.44				6.86				15.0							-	-	21.40	-	-	17.10		-
Manganese	1.90				0.109			0.07		-	-	0.069				0.064				0.127							-	-	0.38	-	-	0.35		-
Nitrate (NO3 as N)	0.7 (7)		0.1	0.02	0.04	0.21	0.07	0.02	< 0.01	< 0.01	0.02	0.09	0.73	5.59	0.6	< 0.01	0.02	0.21	< 0.01	< 0.01	< 0.01	0.06	0.01	0.03	0.02	0.02	0.26	0.03	0.06	0.31	0.39	0.04	0.1	0.04
Nitrite (NO2 as N)	,		< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	0.04	0.05	0.02	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	0.01	0.02	< 0.01	< 0.01	< 0.01	0.07	< 0.01	< 0.01	< 0.01	< 0.01
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	0.62	0.87	0.66	0.26	0.14	0.42	0.29	0.23	0.04	0.35	0.44	7.78	0.93	0.14	0.25	0.11	0.13	0.17	0.32	0.18	0.15	0.15	0.74	20.9	0.38	0.44	0.22	0.91	1.16	0.58	0.64	0.68
Total Kjeldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	1.1	1.9	1.7	0.6	0.50	0.90	0.60	0.80	0.4	0.5	2.2	7.8	1.5	0.4	0.7	0.4	0.5	0.9	0.7	0.7	0.5	0.6	2.1	58.1	1.3	1.5	1.5	3.1	2.60	1.70	2.60	1.60
Dissolved Organic Carbon	1.7	(-/-)			11	-	-	10	-	-		6			-	12	-	-		18				12			-	-	6	-	-	9	-	
Fluoride (Electrode)			-		<0.1	-	-	0.1	-	-	-	0.1			-	0.1	-	-		<0.1				0.10			-	-	0.20	-	-	0.2	-	-
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.14	0.22	0.31	0.18	0.21	0.62	0.16	0.14	0.22	0.19	0.42	0.13	0.15	0.16	0.33	0.27	0.14	0.25	0.21	0.16	0.1	2.55	0.83	2.71	4.24	2.93	1.97	1.56	1.24	3.46	7.98	1.72

Note:

Exceeds ANZECC (2000) guidelines
marine/fresh water ecosystems

Focus of this monitoring report

nc = no comment

NA = no 35

Gerroa GW dw data DP 9-3-16 1/9

Gerroa Waste Dispos	sal Depo	ot																																	
Sample ID	ANZEC	C, 2000	MW4	MW4	MW4	MW4	MW4	MW4	MW4	MW4	MW4	MW4	MW4	MW4	MW4	MW4	MW4	MW4	MW5	MW 5	MW 5	MW5	MW5	MW 5	MW5	MW5	MW5	MW5	MW5	MW5	MW5	MW5	MW5	MW5	MW5
Field Measurements	Fresh	Marine	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16	21/5/10	17/8/10	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	7/2/14
Ground Level (m AHD)			4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.79	4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55
Depth to Groundwater (m AHD)			1.23	0.99	1.04	1.04	1.1	1.43	0.58	0.37	1.04	0.54	0.56	0.69	1.54	1.54	1.34	0.84	0.6	0.81	1.32	1	1.03	1.11	0.72	0.5	0.96	0.85	0.73	0.95	1.24	1.27	0.76	0.56	0.56
Groundwater depth (m bTOC)			4.01	4.25	4.2	4.2	4.14	3.81	4.66	4.87	4.2	4.7	4.68	4.55	3.7	3.7	3.9	4.4	4.15	3.94	3.43	3.75	3.72	3.64	4.03	4.25	3.79	3.9	4.02	3.8	3.51	3.48	3.99	4.19	4.19
Height of Stick up (m)			0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.45	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Groundwater Depth (mbgl)			3.56	3.80	3.75	3.75	3.69	3.36	4.21	4.42	3.75	4.25	4.23	4.1	3.25	3.25	3.45	3.95	3.95	3.74	3.23	3.55	3.52	3.44	3.83	4.05	3.59	3.7	3.82	3.6	3.31	3.28	3,79	3.99	3.99
pH (field)	6.5-8.0 (a)	8-8.4 (a)	6.80	7	6.8	7	6.30	7	7.1	7		7.1			6.6	7	6.4	6.1	7.8	7.7	7.2	7.6	8	8	8	7.9	8.1	7.7	7.7	7	7	7.7	7	7.3	
Temperature (T deg C)	0.0 0.0 (0)	0 0 (2)					-	-	-	-									-	-	-	-	-	-	-	-	-				-	-	-	-	
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		0.58	<1	0.7	0.829	0.44	0.47	0.632	0.683		0.472			555	555	418	207	0.52	0.51	0.414	0.31	0.44	0.31	0.38	0.345	0.36	<1	0.477	0.297	0.13	0.427	0.395	0.425	
Salinity (ppt)						0.020	-	-	-	-									-	-	-	-	-	-	-	-	-			0.20.		-	-	-	
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	3.22	1.1	2.48	1.7	2.26	2.49	2.32	1.97	4.10	2.70	2.10	3.60	3.57	3.39	3.03	2.4	2.81	3.08	2.14	2.22	4.15	2.88	2.15	2.32	2.41	0.9	1.77	1.8	1.52	1.62	2.13	2.09	2.09
Dissolved Oxygen (%)				10.8			-	-												-	22.60	-			-	-	-	9.3						-	
Turbidity (NTU)	6-50 (a)	0.5-10 (a)	-				-	-	-	-									-	-	-	-	-	-	-	-	-				-	-		-	
Redox Potential (mV)				< 0.1				-34				-128				76					-	-		-25				<0.1				-147			
Comments			nc				-			-									nc	nc	nc	nc				-	nc				nc	nc	nc	nc	
Sodium				13				13				10				14					16	-		7				9				15			
Potassium				4			-	4		-		3				2				-	6	-		4		-		5				5			
Calcium			-	117				68				88				90					53			44				60				54			
Magnesium				6			-	6	-	-		5				6			-	-	10	-	-	6	-	-	-	8				7		-	
Chloride				14			-	13	-	-		11				18			-	-	30	-	-	13	-	-	-	15				34		-	
Alkalinity (as CaCO3)			293	323	315	227	137	215	276	284	250	220	258	231	215	230	178	137	142	144	142	120	177	128	147	144	126	179	198	112	172	170	158	155	155
Bicarbonate			293	323	315	227	137	215	276	284	250	220	258	231	215	230	178	137	142	144	142	120	177	128	147	144	121	179	198	112	172	170	158	155	155
Carbonate (as CaCO3)			<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	5	<1	<1	<1	<1	<1	<1	<1	<1
Sulphate (SO4)			-	<1				10				7.00				9					14			4.00				7				4			
pH (lab)							-	-	-	-	6.8	7.1	6.2	6.3					-	-	-	-	-	-	-	-	-				-	-	-	-	
Total Dissolved Solids (TDS)			342	376	406	278	262	248	314	351	304	326	376	311	295	330	258	226	311	238	258	268	252	197	258	204	170	264	326	166	240	224	219	264	264
Hardness (as CaCO3)			-				-	-	-	-									-	-	-	-	-	-	-	-	-				-	-		-	
Total Suspended Solids (TSS)							-		-	-									-	-	-	-	-	-	-	-						-		-	
Iron (filtered)	0.3 (1)			7.98			-	1.61		-		10.3				< 0.05			-	-	< 0.05	-	-	< 0.05	-	-	-	< 0.05			-	3.22		-	
Manganese	1.90		-	0.299			-	0.272		-		0.424				0.086			-	-	0.002	-	-	0.001	-	-		0.001				0.041		-	
Nitrate (NO3 as N)	0.7 (7)		0.07	0.19	0.05	0.15	0.36	0.09	0.03	0.02	0.28	0.03	<0.01	0.19	0.16	0.09	< 0.01	0.02	5.15	10.4	4.44	0.96	0.44	0.14	0.02	1.08	1	1.44	0.03	1.81	< 0.01	0.06	0.14	0.61	0.61
Nitrite (NO2 as N)			<0.01	< 0.01	<0.01	<0.01	0.88	<0.01	< 0.01	< 0.01	< 0.01	<0.01	<0.01	< 0.01	< 0.01	< 0.01	< 0.01	<0.01	0.22	0.21	0.14	0.15	<0.01	0.02	< 0.01	0.12	0.18	0.1	0.08	0.03	0.01	0.01	<0.01	0.11	0.11
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	0.18	0.1	0.15	0.44	1.86	< 0.01	0.37	0.22	0.28	0.09	0.35	0.09	0.75	0.02	0.26	0.12	0.08	< 0.01	0.04	0.03	0.03	0.03	0.08	0.02	0.03	0.05	0.23	0.55	0.45	0.06	5.88	0.64	0.64
Total Kjeldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	0.90	1.2	1.2	0.8	15.30	0.6	1.7	0.7	1.0	0.7	0.8	0.9	2.1	0.3	0.7	0.6	1.7	2.6	2.7	0.6	0.80	0.30	0.70	1.30	1	1.4	2	2.6	17	1.1	9	2.2	2.2
Dissolved Organic Carbon	(-)	(-7	-	7			-	12	-	-		9				3			-	-	6	-	-	5	-	-	-	6				12	-	-	
Fluoride (Electrode)			-	0.2			-	0.2	-	-		0.2				0.20			-	-	0.1	-	-	0.20	-	-	-	0.2			-	0.1		-	
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	1.31	4.29	3.99	0.9	2.4	0.86	2.17	1.6	1.46	1.81	1.56	1.3	1.44	0.06	1.97	1.08	0.09	0.11	< 0.01	0.12	0.1	0.13	0.1	0.08	0.03	0.13	0.22	0.74	1.23	0.33	1.83	1.07	1.07

Note:

Exceeds ANZECC (2000) guidelines
marine/fresh water ecosystems

Focus of this monitoring report
nc = no comment
NA = n 35

Sample ID	ANZEC	2000	MW5	MW5	MW5	MW5	MW5	MW5	MW5	MW5	MW6S	MW 6S	MW 6S	MW6S	MW6S	MW 6S	MW6S	MW6S	MW6S	MW6S	MW6S	MW6S	MW6S	MW6S	MW6S
		.,																							
Field Measurements	Fresh	Marine	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16	21/5/10	17/8/10	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13
Ground Level (m AHD)			4.55	4.55	4.55	4.55	4.55	4.55	4.55	4.55	NA												-	-	<u> </u>
Depth to Groundwater (m AHD)			1.23	0.75	0.77	0.86	1.49	-0.13	1.29	0.85													-	-	-
Groundwater depth (m bTOC)			3.52	4	3.98	3.89	3.26	4.88	3.46	3.9	4.69	4.57	4.45	4.48	4.43	4.39	4.71	4.88	4.5	4.7	4.66	4.43	4.35	4.4	4.77
Height of Stick up (m)			0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Groundwater Depth (mbgl)			3.32	3.8	3.78	3.69	3.06	4.68	3.26	3.7	4.09	3.97	3.85	3.88	3.83	3.79	4.11	4.28	3.9	4.1	4.06	3.83	3.75	3.8	4.17
pH (field)	6.5-8.0 (a)	8-8.4 (a)		6.9			7.5	8	7.8	7	7.5	7.3	6.8	6.9	7.2	7	7.6	7.3	7.5	7.4	7.6	6.9	6.9	7.1	7.4
Temperature (T deg C)											-	-	-										-		-
Electrical Conductivity (mS/cm)	0.125-2.2 (a)			0.126			361	331	410	123	0.52	0.65	0.502	0.37	0.569	0.511	0.504	0.504	0.475	<1	0.396	1.06	0.432	0.474	0.679
Salinity (ppt)											-	-	-	-		-	-	-			0.000		-	-	-
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	2.74	3.2	3.1	2.4					1.74	2.56	2.63	2.19	2.18	2.94	1.9	1.82	2.29	1.2	2.4	2.58	1.46	1.9	2.13
Dissolved Oxygen (%)	0.0 11.0 (u)	0.0 10.0 (u)	2.7 1	32.7	0.1						-	-	27.70	-	-	-	-		-	11.8	2.1	2.00	-	-	-
Turbidity (NTU)	6-50 (a)	0.5-10 (a)									-								-						
Redox Potential (mV)	(-/	0.0 .0 (0)		75.5				56			-	-	-			-33.80		-	-	<0.1			-	-57	-
Comments											nc	nc	nc	nc					nc	nc	nc	nc			-
Sodium				9				12			-		8			44				14				30	-
Potassium				10				2			-		6			8				8			-	6	-
Calcium				16				44			-		80			41				75				75	-
Magnesium				3				5			-	-	9		-	7				6			-	7	-
Chloride				26				18					14			42				21				34	
Alkalinity (as CaCO3)			158	41	118	113	102	118	147	72	182	263	208	176	130	133	196	226	227	202	237	315	62	217	273
Bicarbonate			158	41	118	113	102	118	147	72	182	263	208	176	130	133	196	226	206	202	237	315	62	217	273
Carbonate (as CaCO3)			<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	21	<1	<1	<1	<1	<1	<1
Sulphate (SO4)				<10				5			-	-	26			40.00				18			-	34	-
pH (lab)			7.20	6.90	6.70	6.40																			-
Total Dissolved Solids (TDS)			216	164	234	138	225	199	254	162	314	350	308	332	280	334	328	342	286	314	372	616	286	309	364
Hardness (as CaCO3)											-	-	-										-		-
Total Suspended Solids (TSS)											-	-	-		-								-		-
Iron (filtered)	0.3 (1)			0.25				< 0.05			-		5.59			3.13				< 0.05			-	1.17	-
Manganese	1.90			0.03				< 0.001			-		0.116			0.06				0.025			-	0.019	
Nitrate (NO3 as N)	0.7 (7)		< 0.01	0.63	0.61	< 0.01	0.11	1.4	0.07	0.02	0.37	1.71	0.01	1.16	7.48	0.04	< 0.01	0.8	0.01	0.76	3.54	0.11	< 0.01	1.05	0.62
Nitrite (NO2 as N)			< 0.01	0.03	0.09	< 0.01	<0.01	0.05	0.06	<0.01	0.01	0.02	< 0.01	< 0.01	0.03	< 0.01	<0.01	0.31	0.01	0.08	0.32	< 0.01	<0.01	0.92	0.1
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	4.47	1.28	1.4	14.5	0.03	1.77	0.03	<0.01	0.83	1.8	1.73	0.48	0.23	0.19	0.4	0.25	0.12	0.08	0.2	0.91	0.04	0.29	0.04
Total Kjeldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	6.3	2.5	3.2	18.9	0.6	2.0	0.40	1.0	1.5	2.6	2.5	1	2.40	0.60	0.80	2.00	0.9	3.1	1	4.2	1.9	1.1	1
Dissolved Organic Carbon	\-/	(1)		12				4.00			-	-	8		-	7	-	-	-	4			-	9	-
Fluoride (Electrode)				<0.1				0.20			-	-	0.3			0.60				0.6			-	0.4	-
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	1.03	0.39	0.48	2.08	0.09	1.39	0.02	0.26	0.33	1.04	0.79	0.42	0.68	0.53	3.10	8.81	0.54	0.63	0.4	0.4	0.44	0.12	0.2

Sample ID	ANZEC	0 0000	MANAGO	MW6S	MANAGO	MANCO	MW6S	MW6S	MW6S N	W6S M	W6S MW	B B B B B B	D MM C	D BANAL C	D MW6E	BANKCD	MANA/ CD	MAKED	MAKED	MINICO	MAKED	MAKED	MANAGE	MANAGE	MANAGE	MANAGE	MAKED	MANACO	MW6D	MW6D	MW6D	MANAGE	MW6D	MAKED	MW6D	MW6D
•	-										- 11																									
Field Measurements	Fresh	Marine	7/2/14	7/2/14	-,-,	1/8/14	17/11/14	26/2/15	27/5/15 2	/8/15 4/	1/15 16/2			30/11/1	0 23/2/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	7/2/14	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16
Ground Level (m AHD)			-		NA				NA			N/												-	-	-	-									
Depth to Groundwater (m AHD)			-																					-	-	-	-									
Groundwater depth (m bTOC)			4.94	4.94	4.37	4.77			4.2			4.9	4.82	4.19	4.73	4.7	4.65	4.97	5.12	4.77	4.94	4.91	4.69	4.62	4.65	5.04	5.2	5.2	4.64	5.03	5.01	4.94	4.47	4.63	4.71	5
Height of Stick up (m)			0.6	0.6	0.6	0.6			0.6			0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7
Groundwater Depth (mbgl)			4.34	4.34	3.77	4.17			3.6			4.2	4.12	3.49	4.03	4	3.95	4.27	4.42	4.07	4.24	4.21	3.99	3.92	3.95	4.34	4.5	4.5	3.94	4.33	4.31	4.24	3.77	3.93	4.01	4.3
pH (field)	6.5-8.0 (a)	8-8.4 (a)	7.2			7.3			6.2			7	7	6.6	6.8	7.1	7.2	7.1	6.9	7	7.1	7.2	7.2	7.2	7.2	7.1	7.5			7.1			7	7.2	7	6.8
Temperature (T deg C)	0.0 0.0 (2)	0 011 (0)	-									-	-	-	-	-	-	-		-				-	-	-	-									
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		0.662			0.659			573			1.0	1.02	0.868	0.95	0.996	0.943	1.06	0.994	0.875	<1	0.798	0.86	0.867	0.896	0.851	0.476			0.599			821	801	953	732
Salinity (ppt)			-			0.000						-	-	-	-	-	-	-	-	-			0.00	-	-	-	-			0.000						
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	2.33	2.33	1.85	2			3.48			2.6	2.37	1.34	1.45	2.21	2.71	2.12	1.55	1.45	1	1.81	2.67	1.77	1.58	1.88	1.94	2.75	1.85	1.1	3.1	1.9	2.38	2.51	1.47	2.8
Dissolved Oxygen (%)	()		-			20.9						-	-	14.10		-	-	-	-	-	10.6			-	-	-	-		1100	11.7						
Turbidity (NTU)	6-50 (a)	0.5-10 (a)										-				-			-	-				-												
Redox Potential (mV)			-			65						-	-	-	-	-	30.90	-	-	-	< 0.1			-	13	-	-			43				30		
Comments			-									no	nc	nc	nc					nc	nc	nc	nc	nc	nc	nc	nc									
Sodium			-			24								18			20				15				16					10				13		
Potassium			-			6						-	-	19	-	-	11	-	-	-	10			-	15	-	-			6				6		
Calcium						124						-		120			142				120			-	141					122				139		
Magnesium			-			10						-	-	14	-	-	16	-	-	-	11			-	10	-	-			8				11		
Chloride			-			30						-	-	24	-	-	31	-	-	-	21			-	14	-	-			12				17		
Alkalinity (as CaCO3)			248	<1	182	262			40			54		441	408	396	399	375	410	388	332	342	348	386	375	358	208	325	346	248	271	331	292	305	347	318
Bicarbonate			248	248	182	262						54		441	408	396	399	375	410	388	332	342	348	386	375	358	208	325	346	248	271	331	292	305	347	318
Carbonate (as CaCO3)			<1	<1	<1	<1			<1			<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Sulphate (SO4)			-			60						-		8		-	27	-	-	-	33			-	107	-				71				69		
pH (lab)			-		7	7						-	-	-	-	-	-	-	-	-				-	-	-	-		6.8	7.1	6.7	6.6				
Total Dissolved Solids (TDS)			356	356	280	504			320			47:	504	496	542	534	586	694	636	460	484	508	436	454	496	475	280	438	418	514	497	514	468	522	574	486
Hardness (as CaCO3)			-									-	-	-		-	-	-	-	-				-	-	-	-									
Total Suspended Solids (TSS)			-									-	-	-	-	-	-	-	-	-				-	-	-	-									
Iron (filtered)	0.3 (1)		-			< 0.05						-	-	9.48		-	< 0.05	-	-	-	< 0.05			-	0.07	-	-			< 0.05						
Manganese	1.90		-			0.03						-	-	0.184		-	0.119	-	-	-	0.122			-	0.161	-	-			0.11						
Nitrate (NO3 as N)	0.7 (7)		0.05	0.05	0.06	8.3			<0.01			1.9		0.14	0.17	0.61	5.61	2.93	0.02	0.03	1.29	3.14	0.48	0.41	0.96	0.02	0.02	0.05	0.16	2.92	1.79	0.06	1.13	0.33	<0.01	0.26
Nitrite (NO2 as N)			0.04	0.04	<0.01	0.75			<0.01			<0.0		<0.01		0.02	0.04	0.32	< 0.01	< 0.01	0.08	0.17	0.21	0.03	0.06	0.04	< 0.01	<0.01	0.01	0.08	0.12	0.03	0.06	0.01	<0.01	< 0.01
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	0.04	0.04	0.03	0.02			0.09			16.		21.8	8.9	13.80	3.24	2.25	10.80	7.62	7.06	7.17	12.8	14.60	11.30	10.40	1.97	9.08	8.27	1.79	2.23	3.92	3.13	1.38	4.53	3.48
Total Kjeldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	0.6	0.6	1.5	2.0			2.9			22.	17.5	26.4	16.1	15.20	6.70	9.20	12.40	11.6	9.4	12.4	14	16.5	11.2	10.8	2.2	9.4	8.4	2.8	2.9	4.3	3.20	2.20	6.00	4.20
Dissolved Organic Carbon			-			10							-	15	-	-	11	-			11			-	21	-				11				8.00		
Fluoride (Electrode)			-			0						-	-	0.4	-	-	0.4	-	-	-	0.4			-	0.4	-	-			0.4				0.40		
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.8	0.8	0.2	0.2			0.4			2.9	1	8.06	3.92	4.27	1.39	1.45	6.84	4.80	5.46	0.88	0.38	2.23	2.52	9.07	2.55	3.86	3.93	2.08	0.62	1.74	2.01	1.81	12.00	4.55

Note: Esceeds ANZECC (2000) guidelines 35
Exceeds ANZECC (2000) gu 35

Gerroa Waste Dispos	sal Depo	ot																																			
Sample ID	ANZECO	2000	MW7S	MW 7S	MW 7S	MW7S	MW7S	MW 7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7S	MW7D	MW 7D	MW 7D	MW7D	MW7D	MW 7D	MW7D	MW7D	MW7D	MW7D
Field Measurements	Fresh	Marine	21/5/10	17/8/10	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	7/2/14	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16	21/5/10	17/8/10	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12
Ground Level (m AHD)																												NA				NA					
Depth to Groundwater (m AHD)																																					
Groundwater depth (m bTOC)			4.57	4.44	4.09	4.32	4.31	4.24	4.58	4.76	4.38	4.54	4.56	4.32	4.23	4.26	4.68	4.84	4.84	4.25	4.64	4.67	4.55	4.06	1.9	4.32	4.7	4.76	4.59	4.25	4.46	4.46	4.41	4.72	4.88	4.53	4.7
Height of Stick up (m)			0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.53	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Groundwater Depth (mbgl)			4.04	3.91	3.56	3.79	3.78	3.71	4.05	4.23	3.85	4.01	4.03	3.79	3.7	3.73	4.15	4.31	4.31	3.72	4.11	4.14	4.02	3.53	1.37	3.79	4.17	4.16	3.99	3.65	3.86	3.86	3.81	4.12	4.28	3.93	4.1
pH (field)	6.5-8.0 (a)	8-8.4 (a)	7.5	7.6	7.2	7.4	7.8	7	7.9	7.6	7.9	7.7	7.9	7.5	7.4	7.4	7.8	7.8	1.01	0.72	7.4	1.1.7	1.02	6.8	8.1	7.1	7.3	7.4	7.4	6.9	7.1	7.5	7.4	7.5	7.4	7.5	7.2
Temperature (T deg C)	0.0 0.0 (0)	(-)	-		-			-			-						-	-												-			-			+ :	
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		0.9	1.01	0.836	0.55	0.604	0.392	0.702	0.584	0.522	<1	0.671	0.467	0.442	0.381	0.518	0.58			0.365			317	463	423	579	0.96	0.86	0.712	0.68	0.634	0.649	0.694	0.625	0.661	<1
Salinity (ppt)	0.123-2.2 (a)		-	1.01	-	- 0.55	- 0.004	0.552	0.702	0.504	0.522		0.071	0.407	0.442			-			0.505			317	403	423	313	-	-	0.712	-	0.004	0.043	0.034	- 0.023	0.001	
7 11 7	8.5-11.0 (a)	9.0-10.0 (a)	2.9	2.99	1.36	2.38	2.33	2.83	1.52	2.57	1.81	1.4	2.07	1.65	2.12	1.77	2.03	1.97	1.97	2.15	1.5	2.3	2.7	3.22	3.24	1.82	3.9	1.79	2.59	1.71	2.08	2.53	2.4	1.67	1.61	1.75	0.8
Dissolved Oxygen (%)	0.0 11.0 (u)	0.0 10.0 (4)	-	-	14.30	-	-	-	-	-	-	13.9	2.01	1.00	-	-	-	-	1.01	2.10	15.3	2.0		U.LL	0.21	1.02	0.0	-	-	18.00	-	-	-	-	-	-	7.9
Turbidity (NTU)	6-50 (a)	0.5-10 (a)									-							-														-					
Redox Potential (mV)	0 00 (0)	(-/	-	-	-	-	-	10.50	-	-	-	<0.1				-39		-			83				-75			-	-	-	-	-	23.50	-	-	-	<0.1
Comments			nc	nc	nc	nc					nc				nc	nc	nc	nc										nc	nc	nc	nc		-	-		nc	nc
Sodium				-	109	-		34	-		-	44				40					33				41					24			14	-		-	16
Potassium			-	-	11	-		2			-	6				4		-			6				3				-	16	-		6	-			6
Calcium			-	-	67	-		31			-	56				24		-			56				42				-	90	-		95	-		T . 1	98
Magnesium			-	-	9	-	-	6	-	-	-	7				5		-			7				6			-	-	10	-	-	9	-	-	-	9
Chloride					166			56	-			82				48					59				34					20			14	-		-	41
Alkalinity (as CaCO3)			200	177	204	151	154	85	166	172	147	152	180	132	141	109	153	143	143	143	144	144	159	89	140	133	170	299	348	307	250	226	251	246	262	231	247
Bicarbonate			200	177	204	151	154	85	166	172	136	152	180	132	141	109	153	143	143	143	144	144	159	89	140	133	170	299	348	307	250	226	251	246	262	212	247
Carbonate (as CaCO3)			<1	<1	<1	<1	<1	<1	<1	<1	11	<1	<1	<1	<1	<1	<1	<1	<1	143	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	19	<1
Sulphate (SO4)			-	-	39	-	-	10	-	-	-	14				9		-			8				12			-	-	15	-	-	44	-	-	-	24
pH (lab)																				7	7.4	6.8	6.9					-	-		-			-	-		
Total Dissolved Solids (TDS)			538	528	530	362	256	238	418	328	274	334	398	240	238	201	282	312	312	198	330	399	433	248	267	357	434	449	358	408	408	276	402	432	352	364	402
Hardness (as CaCO3)			-	-		-	-	-	-	-	-							-										-	-	-	-	-	-	-	-	-	
Total Suspended Solids (TSS)			-	-		-					-							-											-				-	-			
Iron (filtered)	0.3 (1)		-	-	0.16	-		1.22				0.4				0.66		-			0.18				0.42				-	5.8			0.07	-			< 0.05
Manganese	1.90		-	-	0.02		-	0.01	-		-	0.014				0.01	-	-			0.013				0.01			-	-	0.106	-	-	0.101	-	-	1	0.081
Nitrate (NO3 as N)	0.7 (7)		0.01	0.01	0.02	0.06	< 0.01	0.04	< 0.01	0.12	<0.01	0.11	< 0.01	0.1	< 0.01	0.16	<0.10	<0.01	<0.01	< 0.01	0.02	< 0.01	< 0.01	0.04	0.05	<0.01	0.02	3.5	0.05	0.12	1.06	0.05	0.16	< 0.01	0.06	0.06	0.47
Nitrite (NO2 as N)	` '		< 0.01	< 0.01	<0.01	< 0.01	< 0.01	< 0.01	< 0.01	0.02	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	0.04	<0.10	0.02	0.02	< 0.01	0.01	0.01	0.02	< 0.01	< 0.01	<0.01	< 0.01	0.04	0.04	0.06	0.05	0.02	<0.01	< 0.01	< 0.01	< 0.01	0.06
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	0.49	1.7	2.56	1.34	1.3	0.16	1.93	2.03	2.25	2.34	2.74	1.2	1.91	0.91	1.33	1.64	1.64	0.68	1.12	0.54	1.16	0.59	0.02	0.8	0.7	21.9	28.6	21.7	10.2	7.99	4.38	4.17	1.98	2.61	1.94
Total Kjeldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	0.8	2.6	4.5	1.6	2.2	0.4	2	3.3	2.6	2.8	3.5	1.4	2.6	1.5	2	1.8	1.8	1.00	1.6	0.7	1.6	0.8	0.8	1.3	1.2	32.7	35.7	28.4	14.5	9.60	4.50	4.80	4.80	3.60	2.40
Dissolved Organic Carbon	\-/	(-/	-	-	9	-	-	6		-	-	6			-	8		-			7				5			-	-	13	-		5	-		$\overline{}$	5
Fluoride (Electrode)			-	-	<0.1	-	-	0.1	-	-	-	0.1			-	<0.1	-				<0.1	1		i e	0.1			-	-	0.1	-	-	0.3	-	T -	1 - 1	0.2
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.25	0.21	0.5	0.74	0.83	0.57	0.32	0.24	0.58	0.32	0.18	0.19	0.16	0.96	0.67	0.22	0.22	0.33	0.23	0.08	0.52	0.21	0.08	0.16	0.32	0.33	0.5	1.52	0.44	0.26	0.61	5.74	3.36	1	0.96

Note:

Exceeds ANZECC (2000) guidelines
marine/fresh water ecosystems

Focus of this monitoring report
nc = no comment
NA = no 35

Gerroa Waste Dispo	sal Depo	ot																					
Sample ID	ANZEC	C, 2000	MW7D	MW7D	MW7D	MW7D	MW7D	MW7D	MW7D	MW7D	MW7D	MW7D	MW7D	MW7D	MW7D	MW7D	MW9	MW 9	MW 9	MW9	MW9	MW 9	MW9
Field Measurements	Fresh	Marine	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16	21/5/10	17/8/10	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11
Ground Level (m AHD)						-	-										1.37	1.37	1.37	1.37	1.37	1.37	1.37
Depth to Groundwater (m AHD)																	0.53	0.55	0.95	0.63	0.52	0.48	0.47
Groundwater depth (m bTOC)			4.67	4.46	4.38	4.4	4.8	4.98	4.39	4.78	4.79	4.69	4.21	4.5	4.45	4.8	1.79	1.77	1.37	1.69	1.8	1.84	1.85
Height of Stick up (m)			0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.95	0.95	0.95	0.95	0.95	0.95	0.95
Groundwater Depth (mbgl)			4.07	3.86	3.78	3.8	4.2	4.38	3.79	4.18	4.19	4.09	3.61	3.9	3.85	4.2	0.84	0.82	0.42	0.74	0.85	0.89	0.9
pH (field)	6.5-8.0 (a)	8-8.4 (a)	7.4	7.5	7.4	7.4	7.4	7.5		7.7			7.4	7.4	7.3	7.3	5.9	6	6.3	6.5	6.7	6.8	6.8
Temperature (T deg C)							-										-		-	-	-	-	
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		0.645	0.622	0.526	0.528	0.547	0.476		0.435			426	483	499	443	21.7	16.6	1.4	8.66	7.39	4.91	6.23
Salinity (ppt)	```				-		-										-			-	-	-	-
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	2.92	2.89	1.54	1.56	2.2	1.94	1.44	1.9	2.9	3.6	3.4	2.06	1.95	3.1	4.74	5.42	3.36	8.68	2.24	2.89	2.14
Dissolved Oxygen (%)					-		-			19.2							-		35.60	-	-	-	
Turbidity (NTU)	6-50 (a)	0.5-10 (a)					-										-				-	-	
Redox Potential (mV)					-	-28	-			73.5				66			-		-	-	-	101	-
Comments			nc	nc	nc	nc	nc	nc									nc	nc	nc	nc	-	-	-
Sodium					-	17	-			12				9			-		759	-	-	904	-
Potassium						6				4				4					33			37	
Calcium						76				70				75					34			31	
Magnesium					-	7	-	-		6				7			-	-	67	-	-	77	-
Chloride					-	16	-	-		8				9			-		912	-	-	1410	-
Alkalinity (as CaCO3)			247	228	203	250	265	208	220	178	212	209	175	214	215	182	85	61	288	129	118	147	116
Bicarbonate			247	228	203	250	265	208	220	178	212	209	175	214	215	182	85	61	288	129	118	147	116
Carbonate (as CaCO3)			<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Sulphate (SO4)					-	18	-	-		8				7			-	-	33	-	-	258	-
pH (lab)					-	-	-	-	7.2	7.7	5.9	6.5					-		-	-	-	-	-
Total Dissolved Solids (TDS)			406	346	338	298	307	280	238	280	292	237	249	300	337	382	15600	10900	3270	6560	3900	2920	3660
Hardness (as CaCO3)					-		-										-		-	-	-	-	-
Total Suspended Solids (TSS)					-	-	-	-									-	-		-	-	-	-
Iron (filtered)	0.3 (1)					0.09				< 0.05				< 0.05					4.64			0.57	-
Manganese	1.90					0.064				0.034				0.037					0.015	-		0.002	-
Nitrate (NO3 as N)	0.7 (7)		0.97	0.46	0.16	0.49	< 0.10	0.02	< 0.01	1.19	0.25	< 0.01	0.46	< 0.01	< 0.01	0.03	0.4	1.98	6.83	0.52	0.19	0.17	< 0.01
Nitrite (NO2 as N)			0.01	0.01	0.01	0.1	<0.10	< 0.01	< 0.01	0.04	0.01	0.02	< 0.01	< 0.01	0.03	< 0.01	< 0.01	<0.01	< 0.50	0.02	<0.01	< 0.01	< 0.01
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	1.67	1.93	2.82	2.43	2.09	1.97	1.91	0.2	1.36	1.48	0.96	0.54	1.64	1.02	<0.10	<0.10	0.84	0.06	0.05	< 0.01	1
Total Kjeldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	2	2.2	3.90	2.60	3	2	2.10	0.60	1.60	1.60	1.30	0.80	2.30	1.40	0.8	2.7	94.1	1.6	2.40	1.70	2.60
Dissolved Organic Carbon					-	13	-	-		7				4			-		430	-	-	51	-
Fluoride (Electrode)					-	0.2	-	-		0.2				0.2			-	-	0.5	-	-	0.4	-
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.08	0.17	1	1.66	8.46	2.55	0.89	0.37	0.30	1.07	0.71	0.19	8.14	1.60	< 0.01	0.83	<2.00	0.07	0.12	0.07	0.22

Note: Exceeds ANZECC (2000) guidelines 35 marine/fresh water ecosystems

Focus of this monitoring report nc = no comment NA = not available 35

Gerroa Waste Dispo	osal Depo	ot																		
Sample ID	ANZEC	C, 2000	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9	MW9
Field Measurements	Fresh	Marine	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	7/2/14	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16
Ground Level (m AHD)			1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37	1.37
Depth to Groundwater (m AHD)			0.39	0.52	0.47	0.88	1.03	0.53	0.4	0.49	0.38	0.38	0.67	0.83	0.54	0.56	0.66	-0.01	0.49	0.52
Groundwater depth (m bTOC)			1.93	1.8	1.85	1.44	1.29	1.79	1.92	1.83	1.94	1.94	1.65	1.49	1.78	1.76	1.66	2.33	1.83	1.8
Height of Stick up (m)			0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95
Groundwater Depth (mbgl)			0.98	0.85	0.9	0.49	0.34	0.84	0.97	0.88	0.99	0.99	0.7	0.54	0.83	0.81	0.71	1.38	0.88	0.85
pH (field)	6.5-8.0 (a)	8-8.4 (a)	6.3	6.7	6.3	6.5	6.7	6.4	6.2	6.4	6.2			5.7			6.4	6.7	6.5	6.3
Temperature (T deg C)			-	-				-	-	-	-									
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		13	6.41	11	9.32	12.7	6.39	1.08	10.2	18.9			23.4			9120	7690	4610	9520
Salinity (ppt)																				
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	2.89	3.31	1.2	1.73	1.68	3.71	1.97	2.14	1.9	1.9	3.69	4.1	2.2	3.3	3.61	1.8	1.92	1.9
Dissolved Oxygen (%)					12.5									45.9						
Turbidity (NTU)	6-50 (a)	0.5-10 (a)																		
Redox Potential (mV)					<0.1				38					243				130		
Comments				nc	nc	nc	nc	nc	nc	nc	nc									
Sodium				-	2330				190					5370				1350		
Potassium				-	101			-	13	-				175				60		
Calcium			-	-	147			-	6	-	-			382				69		
Magnesium				-	392			-	14	-				772				165		
Chloride			-	-	4620			-	275	-	-			8360				1960		
Alkalinity (as CaCO3)			78	174	79	194	191	125	53	130	86	86	97	39	146	135	142	135	133	84
Bicarbonate			78	174	79	194	191	125	53	130	86	86	97	39	146	135	142	135	133	84
Carbonate (as CaCO3)			<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Sulphate (SO4)				-	679			-	37	-				1450				384		
pH (lab)			-	-				-	-	-	-		6.1	5.7	6.1	6.3				
Total Dissolved Solids (TDS)			9530	4000	7670	6540	8220	3320	671	7530	13100	13100	6240	20900	8190	6730	3580	5160	2870	6840
Hardness (as CaCO3)				-				-	-	-										
Total Suspended Solids (TSS)				-				-	-	-										
Iron (filtered)	0.3 (1)			-	0.11			-	0.68	-				0.08				0.26		
Manganese	1.90		-	-	0.005			-	0.004	-	-			0.006				0.004		
Nitrate (NO3 as N)	0.7 (7)		<0.01	0.06	0.11	<0.10	0.02	< 0.01	<0.10	<0.01	0.01	0.01	0.13	1.60	< 0.01	<0.01	<0.01	<0.01	<0.01	0.02
Nitrite (NO2 as N)			<0.01	<0.01	< 0.01	<0.01	<0.01	< 0.01	<0.10	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	<0.10	0.03	0.04	0.23	1.08	< 0.01	0.02	0.32	0.24	0.24	<0.01	<0.01	0.34	0.21	<0.01	0.01	0.47	0.07
Total Kjeldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	1.00	1.9	1.8	2.4	2.8	3.1	3.1	1.6	1.1	1.10	0.40	1.00	2.10	2.60	2.40	4.90	2.20	1.40
Dissolved Organic Carbon				-	17			-	64	-	-			14				0.3		
Fluoride (Electrode)				-	0.2				0.2					0.2				29		
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.08	0.14	0.05	0.17	0.23	0.34	0.24	0.09	0.06	0.06	0.05	0.07	0.22	0.20	0.13	0.31	0.16	0.03

35

Gerroa Waste Dispo	sal Depo	ot																								
Sample ID	ANZEC	C, 2000	MW10	MW 10	MW 10	MW10	MW10	MW 10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10	MW10
Field Measurements	Fresh	Marine	21/5/10	17/8/10	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16
Ground Level (m AHD)			1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475	1.475		1.475	1.475
Depth to Groundwater (m AHD)			0.295	0.295	0.785	0.395	0.295	NA	0.185	0.035	0.275	0.225	0.605	0.765	0.365	0.195	0.235	0.105	0.415	0.475	0.355	0.305	0.365		0.235	0.335
Groundwater depth (m bTOC)			2.14	2.14	1.65	2.04	2.14		2.25	2.4	2.16	2.21	1.83	1.67	2.07	2.24	2.2	2.33	2.02	1.96	2.08	2.13	2.07		2.2	2.1
Height of Stick up (m)			0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96		0.96	0.96
Groundwater Depth (mbgl)			1.18	1.18	0.69	1.08	1.18	NA	1.29	1.44	1.2	1.25	0.87	0.71	1.11	1.28	1.24	1.37	1.06	0.90	1.12	1.17	1.11		1.24	1.14
								INA											1.06		1.12	1.17				
pH (field)	6.5-8.0 (a)	8-8.4 (a)	4	4.1	5.8	5	5.4	-	5.3	4.9	5.2	4.7	5	5.1	5	5.4	5.3	5.1		4.2			5.5		5.6	5.1
Temperature (T deg C)						-	-		-	-						-										└
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		29	6.45	3.06	8.94	5.12	-	5.7	14.8	6.3	12	15.2	13.1	6.64	6.72	14.4	28.2		22.7			8540		4570	8060
Salinity (ppt)					-	-			-	-					-	-										
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	4.69	4.93	4.7	9.61	6.1	-	4.16	3.05	3.26	2.4	2.03	3.43	2.01	3.02	2.38	3.89	3.06	2.7	3.5	4.2	3.29		3.14	2.9
Dissolved Oxygen (%)				-	50.30	-		-	-	-		24.9			-	-	-			30.3						
Turbidity (NTU)	6-50 (a)	0.5-10 (a)					-																			
Redox Potential (mV)			-	-	-	-		-	-	-	-	<0.1			-	161	-	-		318						
Comments			nc	nc	nc	nc			-	-	nc				nc	nc	nc	nc								
Sodium				-	406	-			-	-		2390			-	1100				5060						
Potassium			-	-	20	-			-	-	-	102			-	48	-	-		156						
Calcium					7	-			-	-		114			-	44				265						
Magnesium					13	-			-	-		336			-	136				699						
Chloride				-	449	-		-	-	-		4730			-	1760	-			7760						
Alkalinity (as CaCO3)			<1	<1	28	3	10	-	6	2	6	4	1	<1	5	9	10	4	<1	<1	2	3	16		16	8
Bicarbonate			<1	<1	28	3	10		6	2	6	4	1	<1	5	9	10	4	<1	<1	2	3	16		16	8
Carbonate (as CaCO3)			<1	<1	<1	<1	<1		<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1		<1	<1
Sulphate (SO4)			-	-	42	-		-	-	-	-	642			-	304	-	-		1410						
pH (lab)				-	-	-	-		-	-					-	-			5.20	4.20	5.10	5.10				
Total Dissolved Solids (TDS)			20500	10100	1810	5740	2700	-	3440	10500	3980	8280	10700	7550	3870	3760	10800	19400	7970	16900	12200	8070	3530		3370	6310
Hardness (as CaCO3)			-	-	-	-	-	-	-	-	-				-	-	-	-								
Total Suspended Solids (TSS)																										
Iron (filtered)	0.3 (1)				10.4	-						0.28				0.54				0.27						
Manganese	1.90			-	0.003	-	-		-	-		0.015			-	0.014	-			0.02						
Nitrate (NO3 as N)	0.7 (7)		0.2	0.3	1.36	0.44	0.05	-	0.15	< 0.01	0.03	0.25	< 0.01	0.01	< 0.01	0.04	0.02	0.16	< 0.01	0.72	0.07	0.02	< 0.01		0.05	0.09
Nitrite (NO2 as N)			< 0.01	< 0.01	< 0.50	0.02	< 0.01	-	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	0.02	< 0.01	< 0.01	<0.01	<0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01		< 0.01	< 0.01
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	<0.10	<0.10	<0.10	<0.10	0.03	-	0.02	<0.10	0.02	0.02	<0.10	0.58	0.05	0.06	0.3	0.09	0.08	< 0.01	0.13	0.15	0.05		0.02	< 0.01
Total Kjeldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	2.2	6.2	59.3	1.2	2.6	-	5.2	8	1.5	10	1.4	1.4	3.1	6.1	0.3	<0.2	1.00	2.2	11.4	17.3	3.2		2.5	1.7
Dissolved Organic Carbon		. /		-	430	-	-		-	-	-	12				24	-	-		14						
Fluoride (Electrode)			-	-	<0.1	-	-		-	-		0.2			-	<0.1				0.30						
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.15	0.52	<1.00	0.03	0.14		0.46	0.8	0.3	0.94	0.03	0.04	0.21	0.45	< 0.01	< 0.02	0.09	0.16	1.08	1.67	0.26		0.14	0.06

Note: 25 ANZECC (2000) guidelines 25 marine/fresh water ecosystems

Focus of this monitoring report 10 nc = no comment 10 NA = not available

Gerroa Waste Dispo	sal Depo	ot																								
Sample ID	ANZEC	C, 2000	MW11	MW 11	MW 11	MW11	MW11	MW 11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11	MW11
Field Measurements	Fresh	Marine	21/5/10	17/8/10	30/11/10	23/2/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	6/5/14	1/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16
Ground Level (m AHD)			1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695	1.695
Depth to Groundwater (m AHD)			0.405	0.485	0.955	0.495	0.405	0.375	0.345	0.265	0.375	0.385	0.785	0.965	0.505	0.315	0.375	0.265	0.635	0.705	0.445	0.515	0.575	0.325	0.375	0.355
Groundwater depth (m bTOC)			2.25	2.17	1.7	2.16	2.25	2.28	2.31	2.39	2.28	2.27	1.87	1.69	2.15	2.34	2.28	2.39	2.02	1.95	2.21	2.14	2.08	2.33	2.28	2.3
Height of Stick up (m)			0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96	0.96
					0.74												1.32			0.99						
Groundwater Depth (mbgl)			1.29	1.21		1.2	1.29	1.32	1.35	1.43	1.32	1.31	0.91	0.73	1.19	1.38		1.43	1.06	0.00	1.25	1.18	1.12	1.37	1.32	1.34
pH (field)	6.5-8.0 (a)	8-8.4 (a)	5.1	5.2	5.7	6.1	6.2	6.3	6.4	6	6.1	5.4	5.7	6.2	5.5	6.2	5.7	5.6		5.2			5.9	6.4	6.1	5.6
Temperature (T deg C)				-	-	-		-	-	-	-				-			-								
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		18.6	16.2	0.178	6.12	4.05	1.67	3.83	6.26	3.4	8	12.1	7.05	7.3	1080	4.73	8.28		15.5	4.62	7.85	4200	1730	1590	6370
Salinity (ppt)					-	-			-						-			-								
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	2.41	3.32	5.13	9.66	2.66	1.56	6.61	2.58	2.16	1.2	1.8	1.63	2.05	1.97	2.25	2.15		3.1			2.11	1.8	1.99	2
Dissolved Oxygen (%)			-	-	53.30	-	-	-	-	-	-	12.4			-	-	-	-		33.5						
Turbidity (NTU)	6-50 (a)	0.5-10 (a)		-	-	-			-	-						-		-								
Redox Potential (mV)				-	-	-		111	-	-		<0.1				38		-		206				-30		
Comments			nc	nc	nc	nc			-	-	nc	nc	nc	nc	nc	nc	nc	nc								
Sodium				-	368	-		298	-	-		1900			-	190		-		3710				302		
Potassium				-	16	-		10	-	-		67			-	13	-	-		99				13		
Calcium				-	10	-	-	31	-	-	-	100			-	6	-	-		218				10		
Magnesium					15	-		40	-			249				14		-		582				23		
Chloride			-	-	400	-	-	493	-	-		3690			-	275	-	-		6060						
Alkalinity (as CaCO3)			24	24	51	103	102	93	60	64	79	28	23	87	28	53	32	26	75	7	40	28	68	83	44	34
Bicarbonate			24	24	51	103	102	93	60	64	79	28	23	87	28	53	32	26	75	7	40	28	68	83	44	34
Carbonate (as CaCO3)			<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Sulphate (SO4)				-	30	-	-	121	-	-	-	471			-	37	-	-		1060						
pH (lab)				-	-	-		-	-	-	-				-	-	-	-	6	5	6	6				
Total Dissolved Solids (TDS)			12800	10700	1530	3710	2260	1330	2370	3570	1990	6370	8300	4310	3950	671	2900	5480	1760	13800	2810	4840	2290	1470	1020	4510
Hardness (as CaCO3)				-	-	-			-	-						-	-	-								
Total Suspended Solids (TSS)				-	-	-		-	-	-						-		-								
Iron (filtered)	0.3 (1)			-	6.2	-		2.57	-	-		1.69			-	0.68	-	-		2.98				1.28		
Manganese	1.90		-	-	0.005	-	-	0.01	-	-	-	0.028			-	0.004	-	-		0.03				0.004		
Nitrate (NO3 as N)	0.7 (7)		0.06	0.17	0.93	0.48	<0.01	0.04	<0.01	<0.01	< 0.01	0.08	<0.01	0.81	<0.01	<0.10	<0.01	<0.01	< 0.01	0.09	<0.01	<0.01	<0.01	< 0.01	<0.01	<0.01
Nitrite (NO2 as N)			< 0.01	<0.01	< 0.50	0.01	<0.01	<0.01	<0.01	<0.01	< 0.01	<0.01	<0.01	<0.01	<0.01	<0.10	<0.01	<0.01	< 0.01	<0.01	<0.01	< 0.01	<0.01	< 0.01	<0.01	< 0.01
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	<0.10	<0.10	<0.10	0.13	0.06	0.03	0.03	0.01	0.03	0.05	<0.10	0.42	0.07	0.02	0.01	<0.01	0.06	<0.01	0.19	0.13	<0.01	0.01	0.02	< 0.01
Total Kjeldhal Nitrogen (TKN)	0.5 (5)	0.12 (6)	2.8	3.9	67.7	3.2	4.30	4.70	2.60	3.40	3	5	2.7	2.3	6	3	1.2	5.0	2.5	2.6	3.7	4.9	5.6	4.9	2.2	1.7
Dissolved Organic Carbon					440	-		145	-		-	48				64	-	-		31				102		
Fluoride (Electrode)					0.1			0.2			-	0.2				0.2	-			0.2				0.3		
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.11	0.52	<1.00	0.31	0.25	0.44	0.25	0.12	0.07	0.3	0.06	0.06	0.32	0.2	0.07	0.47	0.14	0.16	0.29	0.32	0.16	0.31	0.07	0.06

Note: Exceeds ANZECC (2000) guidelines marinefresh water ecosystems

Focus of this monitoring report

nc = no comment NA = not available

Sample ID	ANZECO	C, 2000	ML-1	ML-1	ML-1	ML-1	ML-1	ML-1	ML-1	ML-1	ML														
Field Measurements	Fresh	Marine	10/07/03	22/10/03	2/02/04	13/05/04	13/07/04	26/10/04	16/02/05	29/06/05	30/08/05	23/11/05	27/02/06	31/05/06	31/08/06	28/11/06	27/02/07	23/05/07	22/08/07	21/11/07	22/02/08	29/05/08	3 25/08/08	19/11/08	2008
pH (field)	6.5-8.0 (a)	8-8.4 (a)	6.93	7.85	7.58	7.59	7.15	4.31	7.41	6.42	7.13	7.42	7.2	6.45	6.29	6.27	6.9	7.68	5.82	7.09	6.91	6.49	6.5	7.1	No acce
Temperature			12.75	21.66	22.54	17.36	13.33	19.15	18.89	13.65	16.71	19.18	23.57	11.57	15.92	22.4	22.48	11.93	13.53	23.93	20.87	15.26	15.1	20.5	Bros. I
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		1.2	40	13	36.7	13.6	0.4	4.2	4.5	1.941	11.12	36.76	45	9.616	25.73	10.63	10.93	1.388	12.93	1.292	3.591	3.7	9.67	samp 31/0
Eh (ORP) (mV)	0.125-2.2 (a)		-67	nm	nm	nm	nm	nm	nm	-23	nm	nm	nm	nm	nm	nm	nm	76	30/0 27/11						
Salinity (ppt)			0.65	25.53	7.48	23.16	7.93	0.21	2.22	2.4	0.99	6.35	23.27	29.03	5.43	15.73	6.03	6.23	0.7	7.43	0.7	2.37	2.5	nm	
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	4.8	6.2	3.8	2	0.5	0.8	5.1	7.8	8.97	8.26	2.34	5.83	14.29	6.81	4.67	7.88	8.93	5.69	5.36	5.25	7.9	nm	
Dissolved Oxygen (%)	0.0 11.0 (4)	0.0 10.0 (4)	45	82.4	42.6	24.1	5.7	8.3	55.2	64.9	92.8	92.7	31.5	64.4	149.4	86	54.4	74.2	86.1	70.5	60.2	64.9	79	55	
Turbidity (NTU)	6-50 (a)	0.5-10 (a)	12.7	29.3	28	24.8	22.3	13.8	12.4	10.8	10.7	1.9	12.1	17.5	40.3	3.9	15.6	4	4.7	5.3	25.5	3.9	4.9	nm	
Laboratory Analyses	0 00 (0)	1 010 10 (0)																	1	1				1	
Sodium (ICP)			156	9695	2890	8881	2325	41	733	686	396	218	6700	8600	1200	<1	1600	1500	160	2100	150	1200	520	1300	
Potassium (ICP)			8.3	362	122	374	115	8.8	32	33	22	9.4	240	400	83	2.7	76	80	10	96	9.4	53	25	54	
Calcium (ICP)			34	212	106	207	115	13	44	46	45	98	280	150	83	330	65	81	29	81	33	130	50	73	
Magnesium (ICP)			28	-	318	994	305	9.4	93	97	50	28	830	940	160	0.06	200	220	32	280	26	160	63	160	
Chloride							-	-	-	-	-	-	-	-	-		-	-	-	-	-	-	-	-	
Alkalinity (as CaCO3)			235	120	130	119	120	18	75	9.2	74	30	157	100	91	98	53	91	-	120	69	<2	64	90	
Sulphate (SO4)			-	-	-	-	-	-	-	-	-	-	-	-	-	0.013	-	-	-	-	-	-	-	-	
pH			-	-	-	1	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	7	
Redox Potential (mV)			-	-	-	1	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	160	
Total Dissolved Solids (TDS)			12692	32132	6716	19936	8866	220	2304	2404	2032	7664	26108	32000	5600	31000	6600	7500	728	7500	840	2400	2100	6400]
Total Suspended Solids (TSS)			-	47	5	11	14	31	6	18	9	22	22	66	21	93	19	22	11	-	-	-	-	-]
Metals (mg/L)	•																								1
Iron (ICP)	0.3 (1)		0.21	0.12	0.18	<0.002	0.28	3.5	0.072	0.29	0.06	0.33	0.03	2.1	2.7	0.14	0.08	0.41	0.39	0.51	2.5	1.5	0.47	0.7	1
Manganese (ICP)	1.90		0.15	0.028	0.02	0.02	0.027	0.49	0.007	0.29	<0.01	0.06	0.084	0.03	0.1	0.03	0.05	0.1	0.16	0.037	0.14	0.14	0.08	0.06	1
Nutrients (mg/L)		_		,					,	,		,									,				1
Nitrate (NO3 as N)	0.7 (7)		0.03	0.49	0.08	0.19	0.06	0.1	0.04	0.069	<0.04	<0.04	<0.04	0.14	<0.04	0.14	0.16	0.06	0.11	0.07	<0.04	-	-	0.05	1
Nitrite (NO2 as N)			<0.02	0.09	<0.02	0.09	<0.02	<0.02	<0.02	0.016	0.015	0.028	0.022	0.011	0.004	0.06	0.012	0.004	0.03	0.031	0.012	-	-	0.01	1
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	0.05	0.61	1.4	1.5	2.2	<0.02	0.71	<0.02	0.21	0.45	2.1	0.4	1.3	0.49	0.05	0.69	<0.10	1.3	0.31	1.5	0.22	<1	1
Total Kjeldahl Nitrogen (TKN)	0.5 (5)	0.12 (6)	0.38	0.92	1.6	1.6	2.9	1.1	0.94	0.62	0.86	0.94	3.1	1.1	2.3	2.7	0.83	1.4	0.57	1.7	1.3	2.2	0.93	0.8	l
Total Organic Carbon (TOC)			6	1	1	<1	3	27	18	9	7	3	<1	NR	8	<1	5		8	11	16	10	10	-	1
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.02	<0.002	<0.002	0.11	0.05	0.24	0.004	0.023	<0.002	<0.005	0.019	< 0.005	0.015	0.013	0.049	0.005	0.39	0.015	< 0.05	0.01	<0.005	<0.05	Į.
Biological (CFU/100 ml)				ı									1		1			1	7	1	1				
Enterococcus	35 (8)	35 (8)	<1	12	400	4 (app)	40	90	90	<1	38	120	36	30 (app)	20 (app)	20 (app)	1500 (app)	72 (app)	<2	20	102	33	2 (app)	18 (app)	
Thermotolerant (Faecal) coliforms	150 (8)	150 (8)	<1	4	14 (app)	6 (app)	12 (app)	250	20 (app)	2	72	30	16	12 (app)	20 (app)	<2	1800(app)	12 (app)	<2	128	24	23	10	20	

Exceeds ANZECC (2000) guidelines

Focus of this monitoring report

- 1. Trigger value is an indicative interim working level only (IIWL).
- 2. Ammonia trigger value specified at pH = 8.0, for a 95% protection level, is corrected for an average pH = 7.3. 7. Trigger value for a 95% Protection level
- 3. Trigger value for oxides of Nitrogen (NOx) for lowland rivers in NSW.
- 4. Trigger value for oxides of Nitrogen (NOx) for marine ecosystems in NSW.
- Trigger value for total Nitrogen in lowland rivers in NSW.

- 6. Trigger value for total Nitrogen in marine ecosystems in NSW.
- 8. Guidelinefor water quality and aesthetics: Primary Contact
- a. Reference only, not dirtectly applicable to groundwater.

Sample ID	ANZECO	C, 2000	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2														
Field Measurements	Fresh	Marine	10/07/03	22/10/03	2/02/04	13/05/04	13/07/04	26/10/04	16/02/05	29/06/05	30/08/05	23/11/05	27/02/06	31/05/06	31/08/06	28/11/06	27/02/07	23/05/07	22/08/07	21/11/07	22/02/08	29/05/08	25/08/08	19/11/08	18/02/09
pH (field)	6.5-8.0 (a)	8-8.4 (a)	7.4	8.27	7.75	7.59	7.46	6.72	7.39	6.69	6.74	7.3	7.24	6.45	6.25	5.67	6.13	7.62	6.19	7.27	6.85	6.02	6.7	7.23	7
Temperature			13.71	22.37	24.3	16.68	14.71	19.04	20.67	13.71	17.33	21.78	21.25	11.79	15.84	23.31	23.13	14.46	13.97	25.39	22.11	15.44	14.1	21	-
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		5	41.4	27.4	36.9	23.6	1.3	25.6	6.3	8.986	41.19	47.28	33.07	15.21	47.9	17.5	36.12	2.171	26.73	35.684	14.356	8.5	41.6	20
Eh (ORP) (mV)			-40	nm	nm	nm	nm	nm	nm	-431	nm	nm	nm	nm	nm	nm	nm	120	-						
Salinity (ppt)			2.67	26.24	16.77	23.38	14.25	0.66	15.58	3.42	5.05	26.42	30.81	20.68	8.9	31.24	10.33	22.85	1.12	16.36	23.99	10.4	6.1	nm	-
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	4.5	5.4	3.5	2	0.4	1.5	4.2	6.5	7.6	6.09	3.26	9.5	15.46	6.15	4.64	4.52	7.8	9.81	5.56	6.14	93	nm	8.1
Dissolved Oxygen (%)			44.2	72.7	45.4	23.1	4.1	15.9	51.5	62	81.6	80.8	44.1	100	164.8	86.3	54.6	44.3	76.2	131.2	73.2	75.8	9.5	32	-
Turbidity (NTU)	6-50 (a)	0.5-10 (a)	17.9	40	42.4	26.6	26.3	15.7	18.3	12.1	9.8	4.6	2.1	3.7	44.7	6.8	9.7	4.9	6.1	16.2	3.8	4.2	5.7	nm	-
Laboratory Analyses																									
Sodium (ICP)			742	9217	5760	9283	4430	177	4720	1007	1295	8304	8800	6300	1500	<1	3000	5600	320	4200	6500	5600	1300	7200	3800
Potassium (ICP)			28	345	224	388	181	14	287	48	65	349	330	280	150	0.71	120	260	17	180	270	230	55	290	160
Calcium (ICP)			59	212	181	203	153	20	293	55	66	217	360	120	100	-	95	150	32	140	170	270	70	200	170
Magnesium (ICP)			95	-	765	106	536	26	820	136	162	1015	1100	660	250	0.021	320	800	47	530	790	680	140	890	480
Chloride			-	-	-	-	-	-	-	-	-	-	-	-	-	350	-	-	-	-	-	-	-	-	-
Alkalinity (as CaCO3)			306	125	130	13	135	25	118	11.5	106	109	124	90	120	111	63	120	-	110	110	<2	71	120	25
Sulphate (SO4)			-	-	-	-	-	-	-	-	-	-	-	-	-	0.021	-	-	-	-	-	-	-	-	-
рН			6.5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	7.3	-
Redox Potential (mV)			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	180	-11.4
Total Dissolved Solids (TDS)			3392	32400	20008	21828	16284	672	15624	3612	5540	27196	36608	22000	9000	34000	11000	27000	1050	16000	23000	9600	5000	28000	13000
Total Suspended Solids (TSS)			-	58	28	11	23	27	25	12	21	66	23	50	48	94	16	66	3	-	-	-	-	-	-
Metals (mg/L)																-					_				
Iron (ICP)	0.3 (1)		0.42	0.092	0.48	<0.002	0.21	3.2	0.016	0.1	0.06	0.21	0.01	0.1	2.4	<0.04	0.07	0.28	0.7	0.45	0.26	1.5	0.57	0.34	0.52
Manganese (ICP)	1.90		0.12	0.022	0.026	0.02	0.025	0.45	0.025	0.28	0.01	0.03	0.033	0.02	<0.1	0.03	0.05	0.11	0.15	0.03	0.038	0.04	0.06	0.03	0.13
Nutrients (mg/L)																									
Nitrate (NO3 as N)	0.7 (7)		0.06	<0.02	0.03	0.28	< 0.02	0.03	0.06	0.076	0.08	<0.04	<0.04	0.13	<0.04	350	0.34	0.08	0.08	0.1	0.02	-	-	0.02	0.69
Nitrite (NO2 as N)			<0.02	<0.02	<0.02	0.07	<0.02	<0.02	0.02	0.014	0.059	0.018	0.009	0.011	0.013	8500	0.012	0.017	0.04	0.057	0.025	-	-	0.01	0.2
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	2.3	0.08	1.4	0.95	3.3	0.2	2.1	0.05	3.5	0.32	0.6	0.22	3.8	0.2	0.12	0.96	0.24	1.1	0.88	3.7	0.44	<1	1.9
Total Kjeldahl Nitrogen (TKN)	0.5 (5)	0.12 (6)	2.4	0.08	1.4	1.3	4.1	2	2.1	0.73	4.3	0.73	0.73	0.58	5.2	0.71	0.85	1.4	1	2	1	4.3	0.98	0.7	1.9
Total Organic Carbon (TOC)			5	1	<1	<1	1	26	13	9	7	<1	<1	NR	5	<1	3		8	11	3	5	7	-	-
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.008	<0.002	0.14	0.12	2.8	0.25	0.004	0.021	<0.002	0.007	0.055	0.076	0.014	0.021	<0.005	0.016	0.54	0.15	0.06	0.01	<0.005	0.08	0.01
Biological (CFU/100 ml)																							_		
Enterococcus	35 (8)	35 (8)	7	<1	500(app)	10 (app)	24 (app)	100	50	<1	47	120	48	690	8 (app)	32 (app)	1600 (app)	18 (app)	<2	24	80	35	16 (app)	24 (app)	130
Thermotolerant (Faecal) coliforms	150 (8)	150 (8)	15	<1	72	8 (app)	24 (app)	200	60	2	32	80	50	1000	48	6 (app)	1600 (app)	36 (app)	<2	140	110	17	18	38	120

Exceeds ANZECC (2000) guidelines

0.054

Focus of this monitoring report

Table SW-1: Summary Analytical Results for Surface Water Locations (2003 to 2016)

Sample ID	ANZECO	c, 2000	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2	ML-2
Field Measurements	Fresh	Marine	19/05/09	27/08/09	27/11/09	26/2/10	21/05/10	17/08/10	30/11/10	23/02/11	24/5/11	24/8/11	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	6/5/14	18/8/14	17/11/14	26/2/15
pH (field)	6.5-8.0 (a)	8-8.4 (a)	6.8	6.3	6.7	7.1	6.8	6.7	5.7	6.8	7.2	6.8	6.9	7	7.2	7	7.3	7.4	6.5	7	7.2	6.9	7.1	7	7	6.5
Temperature			-	-	-	-	-	-	-	-									-	-	-	-				
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		33	38	24	21.2	28.9	10.4	1.37	20.1	20	1.41	6.73	19	15.7	20	46.9	19.6	5.13	6.88	13.6	40	38200	21000	36600	553
Eh (ORP) (mV)			-	-	-	-	-	-	-	-									-	-	-	-		149		
Salinity (ppt)			-	-	-	-	-	-	-	-									-	-	-	-				
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	7.2	7.5	6.6	5.2	7	9.92	3.64	5.36	8.2	6.05	6.68	6.25	5.66	3.2	6.03	5.39	4.61	4.97	11.1	5.22	6.66	6.2	5.8	4.1
Dissolved Oxygen (%)			-	89	-	-	-	-	39.10	-						33.2			-	-	-	-		67.3		
Turbidity (NTU)	6-50 (a)	0.5-10 (a)	-	-	-	-	-	-	-	-									-	-	-	-				
Laboratory Analyses																										
Sodium (ICP)			-	8000	-	-	-	-	40	-		189				7180			-	1200	-	-		5400		
Potassium (ICP)			-	390	-	-	-	-	4	-		10				272			-	52	-	-		246		
Calcium (ICP)			-	320	-	-	-	-	7	-		32				255			-	79	-	-		241		
Magnesium (ICP)			-	1000	-	-	-	-	6	-		29				701			-	140	-	-		614		
Chloride			-	-	-	-	-	-	-	-									-	-	-	-				
Alkalinity (as CaCO3)			110	120	110	117	109	64	16	123	114	50	66	128		121	128	117	43	160	127	115	124	92	111	17
Sulphate (SO4)			-	-	-	-	-	-	-	-									-	-	-	-				
рН			-	-	-	-	-	-	-	-									-	-	-	-				
Redox Potential (mV)			-	88	-	-	-	-	-	-		73.5				<0.1			-	58	-	-				
Total Dissolved Solids (TDS)			23000	29000	20000	14200	23500	12000	116	17900	13200	846	6730	13700	11800	18100	34600	12900	2950	4260	9920	28800	32600	18600	24400	423
Total Suspended Solids (TSS)			-	-	-	-	-	-	-	-									-	-	-	-				
Metals (mg/L)		ı			ı		I	ı		T		1	T.	1	1	ı		I			T	T			1	т
Iron (ICP)	0.3 (1)		-	<0.01	-	-	-	-	1.19	-		0.89				<0.50			-	1.45	-	-		0.36		
Manganese (ICP)	1.90		-	<0.01	-	-	-	-	0.057	-		0.12				0.031			-	0.052	-	-		0.025		<u> </u>
Nutrients (mg/L)									,									,			,	,				
Nitrate (NO3 as N)	0.7 (7)		0.1	0.1	0.11	0.05	0.16	0.1	0.05	0.13	0.15	0.08	0.12	0.07	0.24	0.18	<0.01	0.23	0.02	0.15	0.06	0.01	0.03	0.15	<0.01	0.05
Nitrite (NO2 as N)			0.02	0.02	0.04	0.02	0.08	0.01	<0.01	0.16	0.06	<0.01	0.02	0.09	0.06	0.02	<0.01	<0.01	0.02	0.02	0.06	0.05	0.02	0.04	0.04	<0.01
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	0.56	0.41	1.5	1.9	<0.10	<0.10	0.08	0.87	2.39	1.69	1.03	1.7	1.8	1.09	<0.10	<0.10	0.75	3.19	1	0.44	0.33	0.68	0.36	0.09
Total Kjeldahl Nitrogen (TKN)	0.5 (5)	0.12 (6)	0.81	0.41	1.9	4.9	1.1	1.4	1.6	2	3.2	2	1.6	2.4	2.9	1.3	0.2	1	2	3.6	2.2	1	<0.5	1.2	0.9	1.4
Total Organic Carbon (TOC)			-	-	-	-	-	-	-	-									-	-	-	-				
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.01	0.009	1.1	0.1	<0.01	0.02	0.1	<0.01	0.04	0.1	0.03	0.06	0.05	<0.01	<0.01	0.13	0.09	0.04	0.01	0.12	0.6	0.05	0.09	0.21
Biological (CFU/100 ml)		ı					r	ı		r			ı		1			r			ı	ı			1	
Enterococcus	35 (8)	35 (8)	-	-	-	-	-	-	~1300	-		~18				4			-	20	-	-		160		
Thermotolerant (Faecal) coliforms	150 (8)	150 (8)	-	9	-	-	-	-	~6600	-		~17				2			-	24	-	-		290		

Exceeds ANZECC (2000) guidelines

0.054

Focus of this monitoring report

- Trigger value is an indicative interim working level only (IIWL).
 Ammonia trigger value at pH =8, 95% protection, corrected ave pH=7.3.
 Trigger value for oxides of Nitrogen (NOx) for lowland rivers in NSW.
 Trigger value for oxides of Nitrogen (NOx) for marine ecosystems in NSW.
 Reference only, not dirtectly applied.

Sample ID	ANZECO	C, 2000	ML-2	ML-2	ML-2	ML-2	ML-3	ML-3	ML-3	ML-3	ML-3	ML-3	ML-3	ML-4							
Field Measurements	Fresh	Marine	27/5/15	24/8/15	4/11/15	16/2/16	22/10/03	2/02/04	13/05/04	13/07/04	26/10/04	2004-2014	2015-2016	10/07/03	22/10/03	2/02/04	13/05/04	13/07/04	26/10/04	16/02/05	29/06/05
pH (field)	6.5-8.0 (a)	8-8.4 (a)	6.6	7.5	7.1	6.5	7.63	7.42	7.65	7.26	4.25	No access	No access (Cleary	6.99	7.66	7.77	7.7	17.24	4.26	7.33	6.48
Temperature							20.5	21.34	13.03	13.16	18.75	(Cleary Bros. Land) - no	Bros. Land) - no samples taken	12.7	19.99	21.72	12.68	12.43	18.66	19	13.77
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		4280	7790	9400	25400	13.1	5.3	3.1	8.5	0.4	samples taken 18/02/2009;	31/05/2013; 30/08/2013;	1.2	13.3	5.4	3.2	8.1	0.4	2.7	4.4
Eh (ORP) (mV)							nm	nm	nm	nm	nm	19/05/2009;	27/11/2013 an	-67	nm	nm	nm	nm	nm	nm	-172
Salinity (ppt)							7.53	2.85	1.62	4.72	0.21	27/08/2009; 2		0.63	7.63	2.93	1.7	4.48	0.21	1.44	2.33
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	4.46	9.11	4.93	5.4	4.6	4.5	2.2	0.8	1.2			5.1	4	4.1	2.5	0.6	0.5	4.6	7.4
Dissolved Oxygen (%)							52.9	41.1	24	7.6	12.2			46.5	46.1	47.6	23.3	6.6	4.9	50.3	71.6
Turbidity (NTU)	6-50 (a)	0.5-10 (a)					30.1	27.1	21.5	20.6	15.1			17.3	28.8	28.4	22.7	20.4	14.7	26	32.8
Laboratory Analyses																					
Sodium (ICP)				1420			-	-	-	-	-			152	2612	1080	584	1589	40	461	677
Potassium (ICP)				55			-	-	-	-	-			8	103	48	26	57	9.3	23	33
Calcium (ICP)				66			-	-	-	-	-			34	121	62	50	90	13	38	47
Magnesium (ICP)				158			-	-	-	-	-			28	-	119	76	174	9.2	61	95
Chloride							-	-	-	-	-			-	-	-	-	-	-	-	-
Alkalinity (as CaCO3)			68	103	131	102	-	-	-	-	-			282	93	108	83	95	14	60	11.5
Sulphate (SO4)							-	-	-	-	-			-	-	-	-	-	-	-	-
рН							-	-	-	-	-			6.5	-	-	-	-	-	-	-
Redox Potential (mV)				-17			-	-	-	-	-			-	-	-	-	-	-	-	-
Total Dissolved Solids (TDS)			2060	4840	6240	17200	-	-	-	-	-			744	8944	3480	1892	5176	160	1532	2496
Total Suspended Solids (TSS)							-	-	-	-	-			-	48	-	2	7	40	15	36
Metals (mg/L)											ı						1				
Iron (ICP)	0.3 (1)			1.64			-	-	-	-	-			0.23	2.9	0.39	0.27	0.2	3.4	0.095	0.1
Manganese (ICP)	1.90			0.023			-	-	-	-	-			0.15	0.096	0.007	0.04	0.032	0.48	<0.001	0.29
Nutrients (mg/L)											ı				I		ı				I
Nitrate (NO3 as N)	0.7 (7)		0.02	0.13	0.05	0.02	-	-	-	-	-			0.04	0.03	0.12	<0.02	0.08	0.04	0.02	0.075
Nitrite (NO2 as N)			0.02	0.02	0.05	0.02	-	-	-	-	-			<0.02	<0.02	<0.02	0.02	<0.02	<0.02	<0.02	0.019
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	0.8	1.5	1.76	0.81	0.08	0.04	0.08	0.07	0.03			0.04	0.02	<0.02	<0.02	0.06	<0.02	0.21	<0.02
Total Kjeldahl Nitrogen (TKN)	0.5 (5)	0.12 (6)	1.2	1.9	3	1.6	-	-	-	-	-			0.37	0.15	0.75	0.48	0.7	2.1	0.8	0.66
Total Organic Carbon (TOC)							-	-	-	-	-			7	5	3	8	6	28	10	9
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.16	0.1	0.05	0.03	-	-	-	-	-			<0.002	0.049	0.036	<0.002	<0.002	0.3	0.021	0.04
Biological (CFU/100 ml)								1									1		1		
Enterococcus	35 (8)	35 (8)		590			10	390	72	60	140			<1	10	320	60	72	100	180	<1
Thermotolerant (Faecal) coliforms	150 (8)	150 (8)		640			16	32 (app)	50	18 (app)	220			<1	15	28 (app)	44	20 (app)	240	24 (app)	<1

Exceeds ANZECC (2000) guidelines

marine ecosystems in NSW.

Focus of this monitoring report

ı level.

nm = not measured (app) = approximately NR = no result

sthetics: primary contact. cable to groundwater.

Table SW-1: Summary Analytical Results for Surface Water Locations (2003 to 2016)

Sample ID	ANZECO	c, 2000	ML-4	ML-4														
Field Measurements	Fresh	Marine	30/08/05	23/11/05	27/02/06	31/05/06	31/08/06	28/11/06	27/02/07	23/05/07	22/08/07	21/11/07	22/02/08	29/05/08	25/08/08	19/11/08	2008-2014	2015-2016
pH (field)	6.5-8.0 (a)	8-8.4 (a)	7.22	7.67	7.43	6.58	6.42	6.04	6.76	7.6	5.86	7.19	6.94	6.64	6.8	7.07	No access (Cleary Bros.	No access
Temperature			16.62	18.86	22.14	12.58	15.36	20.6	22.61	14.2	13.56	23.26	20.85	14.86	14.3	20	Land) - no samples taken 18/02/2009;	
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		1.98	5.547	10.24	15.62	5.126	13.68	10.9	24.37	1.381	9.749	1.825	1.533	1.7	5.49	19/05/2009; 27/08/2009	
Eh (ORP) (mV)			nm	89	2													
Salinity (ppt)			1.01	3.01	5.79	9.15	2.77	7.92	6.19	14.84	0.7	5.49	1.01	0.97	1.1	nm		
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	9.48	8.26	5.06	6.3	14.45	7.36	5.02	4.71	8.19	5.31	4.88	5.11	8.2	nm		
Dissolved Oxygen (%)			97.9	90.4	60	62.7	146.8	85.8	58.3	46.2	79	64.2	54.9	63.2	81	46		
Turbidity (NTU)	6-50 (a)	0.5-10 (a)	10.5	1	3.2	20.2	31.6	2.8	14.4	5.6	4.7	0	30.5	1.1	4.6	nm		
Laboratory Analyses									•	•						•		
Sodium (ICP)			219	907	3600	2500	670	3	1700	3600	150	1500	230	440	200	740		
Potassium (ICP)			13	38	230	130	39	0.75	75	160	10	70	14	20	11	30		
Calcium (ICP)			39	66	180	67	59	160	66	120	29	70	36	100	40	67		
Magnesium (ICP)			34	122	450	290	85	0.008	210	500	32	210	36	64	34	96		
Chloride			-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Alkalinity (as CaCO3)			71	49	119	15	63	83	52	100	-	90	97	<2	61	80		
Sulphate (SO4)			-	-	-	-	-	0.017	-	-	-	-	-	-	-	-		
рН			-	-	-	-	-	-	-	-	-	-	-	-	-	7.2		
Redox Potential (mV)			-	-	-	-	-	-	-	-	-	-	-	-	-	170		
Total Dissolved Solids (TDS)			1232	4236	13124	9400	3100	17000	6700	17400	602	6400	1000	1100	990	3100		
Total Suspended Solids (TSS)			8	12	9	56	18	43	16	13	15	-	-	-	-	-		
Metals (mg/L)																		
Iron (ICP)	0.3 (1)		0.05	0.28	0.05	3.2	2.4	<0.04	0.06	0.36	0.4	0.22	2.9	0.68	0.43	1.3		
Manganese (ICP)	1.90		<0.01	0.06	0.053	0.1	0.1	0.03	0.05	0.06	0.16	0.017	0.12	0.04	0.09	0.08		
Nutrients (mg/L)																		
Nitrate (NO3 as N)	0.7 (7)		<0.04	<0.04	<0.04	1.4	<0.04	190	0.16	0.09	0.1	0.05	<0.04	-	-	0.05		
Nitrite (NO2 as N)			0.01	0.003	0.008	0.016	0.003	4600	0.012	0.007	0.02	0.011	0.01	-	-	<0.01		
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	<0.10	0.05	0.55	0.29	0.37	<0.1	0.09	1.5	0.37	<0.02	1.7	<0.10	0.27	<1		
Total Kjeldahl Nitrogen (TKN)	0.5 (5)	0.12 (6)	0.62	0.42	1.3	2	1.4	0.75	0.83	2.3	0.44	0.49	2.9	0.53	0.76	0.8		
Total Organic Carbon (TOC)			10	6	2	NR	10	3	6		8	11	16	10	10	_		
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	<0.002	<0.005	0.029	0.098	0.01	0.017	0.036	0.008	0.44	0.019	0.05	<0.005	<0.005	<0.05		
Biological (CFU/100 ml)																	1	
Enterococcus	35 (8)	35 (8)	52	160	100	80	60	30 (app)	1800(ap)	230	<2	32	70	18	8 (app)	14 (app)		
Thermotolerant (Faecal) coliforms	150 (8)	150 (8)	64	25	60	112	30 (app)	14 (app)	2000	12 (app)	<2	16 (app)	16 (app)	15	13	16 (app)		

Exceeds ANZECC (2000) guidelines 0.054

Focus of this monitoring report

Sample ID	ANZECO	C, 2000	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5
Field Measurements	Fresh	Marine	10/07/03	22/10/03	2/02/04	13/05/04	13/07/04	26/10/04	19/11/08	18/02/09	19/05/09	27/08/09	27/11/09	26/2/10	21/05/10	17/08/10	30/11/10	23/02/11	24/5/11	24/8/11
pH (field)	6.5-8.0 (a)	8-8.4 (a)	7.26	7.97	7.54	7.51	7.51	6.6	7.23	6.9	6.7	5.9	6.3	6.8	6.7	6.7	5.8	6.8	7	6.8
Temperature			13.6	22.51	23.47	16.41	14.26	19.03	20.2	-	-	-	-	-	-	-	-	-		
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		3	40.3	19.7	36.1	14.6	0.7	16.67	13	27	31	19	9.6	21.3	5.1	0.294	16.6	8	9.19
Eh (ORP) (mV)			-63	nm	nm	nm	nm	nm	88	-	-	-	-	-	-	-	-	-		
Salinity (ppt)			1.58	25.7	11.7	22.78	8.48	0.34	nm	-	-	-	-	-	-	-	-	-		
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9.0-10.0 (a)	4.3	5.9	3	1.6	0.5	1	nm	8.1	6.4	5.4	4.9	5.2	6.36	9.44	5.18	6.24	5.38	6.39
Dissolved Oxygen (%)			41.8	79	37	18.2	4.8	11.3	67	-	-	62	-	-	-	-	54.00	-		
Turbidity (NTU)	6-50 (a)	0.5-10 (a)	15.4	33.3	26.8	26.1	32.3	14.1	nm	-	-	-	-	-	-	-	-	-		
Laboratory Analyses																				
Sodium (ICP)			-	-	-	-	-	-	2100	2100	-	6300	-	-	-	-	21	-		102
Potassium (ICP)			-	-	-	-	-	-	84	88	-	310	-	-	-	-	3	-		7
Calcium (ICP)			-	-	-	-	-	-	90	110	-	200	-	-	-	-	7	-		31
Magnesium (ICP)			-	-	-	-	-	-	240	260	-	820	-	-	-	-	5	-		21
Chloride			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Alkalinity (as CaCO3)			-	-	-	-	-	-	110	110	110	120	100	106	103	55	6	129	112	45
Sulphate (SO4)			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
pH			-	-	-	-	-	-	7	-	-	-	-	-	-	-	-	-		
Redox Potential (mV)			-	-	-	-	-	-	190	-4	-	82	-	-	-	-	-	-		72.6
Total Dissolved Solids (TDS)			-	-	-	-	-	-	1200	8800	24000	24000	12000	6380	15400	3060	146	10800	4450	538
Total Suspended Solids (TSS)			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Metals (mg/L)	•																			
Iron (ICP)	0.3 (1)		-	-	-	-	-	-	0.57	0.66	-	0.11	-	-	-	-	0.98	-		0.89
Manganese (ICP)	1.90		-	-	-	-	-	-	0.06	0.26	-	<0.01	-	-	-	-	0.062	-		0.139
Nutrients (mg/L)	•																			
Nitrate (NO3 as N)	0.7 (7)		-	-	-	-	-	-	0.08	1.4	0.1	0.2	0.13	0.19	0.16	0.08	0.03	0.36	0.1	0.05
Nitrite (NO2 as N)			-	-	-	-	-	-	0.02	0.43	0.02	0.09	0.04	0.02	0.1	<0.01	<0.01	0.1	0.03	<0.01
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	1.9	1.2	4.1	1.4	4.2	0.2	1.6	2	1.2	2.8	3	2.17	1.52	0.94	0.04	1.83	3.12	1.6
Total Kjeldahl Nitrogen (TKN)	0.5 (5)	0.12 (6)	-	-	-	-	-	-	2.4	2.1	1.2	2.9	3.6	4.6	1.7	1.6	1.5	3.3	4.2	2
Total Organic Carbon (TOC)			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	-	-	-	-	-	-	<0.05	<0.005	0.01	0.04	<0.05	1.78	<0.01	<0.01	<0.01	<0.01	0.03	0.12
Biological (CFU/100 ml)																				
Enterococcus	35 (8)	35 (8)	2	4	600	36 (app)	40	80	116 (app)	180	-	-	-	-	-		~1200	-		~8
Thermotolerant (Faecal) coliforms	150 (8)	150 (8)	<1	16	44	12 (app)	12 (app)	160	58	64	-	6	-	-	-	-	~9100	-		<2

Exceeds ANZECC (2000) guidelines

Focus of this monitoring report

Table SW-1: Summary Analytical Results for Surface Water Locations (2003 to 2016)

Sample ID	ANZECO	2000	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5	ML-5
Field Measurements	Fresh	Marine	3/11/11	1/2/12	31/5/12	10/8/12	21/11/12	18/2/13	31/5/13	30/8/13	27/11/13	7/2/14	6/5/14	18/8/14	17/11/14	26/2/15	27/5/15	24/8/15	4/11/15	16/2/16
pH (field)	6.5-8.0 (a)	8-8.4 (a)	6.9	7	7.2	6.9	7	7.3	6.4	7	7.2	172111	6.5	6.7	6.4	6.1	6.9	7.3	7.1	6.8
Temperature		()										_							!	
Electrical Conductivity (mS/cm)	0.125-2.2 (a)		2.89	9.59	5.08	5	45.9	18.9	1.26	1.81	4.26		26800	12900	26800	357	1210	2380	1820	9900
Eh (ORP) (mV)	0.125-2.2 (a)		2.03	3.33	3.00		43.3	10.5	-	1.01	-	_	20000	208	20000	337	1210	2300	1020	3300
Salinity (ppt)														200						
Dissolved Oxygen (mg/L)	8.5-11.0 (a)	9 0-10 0 (a)	6.86	6.68	7.37	3.6	6.82	4.5	5.22	3.91	6.65	_	5.55	6.2	5.6	5.3	5.2	7.34	3.83	5.1
Dissolved Oxygen (%)	0.3-11.0 (a)	3.0-10.0 (a)	0.00	0.00	1.51	33.7	0.02	4.5	5.22	5.51	0.00	_	3.33	63.3	3.0	3.3	5.2	7.54	3.03	5.1
Turbidity (NTU)	6-50 (a)	0.5-10 (a)				33.1				-				00.0						
Laboratory Analyses	0 00 (u)	0.0 10 (u)																		
Sodium (ICP)						1030				274				2980				379		
Potassium (ICP)						43			-	16	_	_		138				19		
Calcium (ICP)						74			-	48	_	_		154				42		
Magnesium (ICP)						122			-	36	-	-		363				49		
Chloride									-	_	-	-								
Alkalinity (as CaCO3)			52	148	132	111	117	114	37	148	112		138	105	157	15	60	134	134	102
Sulphate (SO4)									-	-	-	-								
pH									-	-	-	-								
Redox Potential (mV)						<0.1			-	62	-	-						95		
Total Dissolved Solids (TDS)			1660	6530	2880	3890	29700	12700	692	936	2510		21400	8160	18500	281	538	1370	1270	6860
Total Suspended Solids (TSS)									-	-	-	-								
Metals (mg/L)																				
Iron (ICP)	0.3 (1)					0.94			-	1.99	-	-		0.6				2.09		
Manganese (ICP)	1.90					0.042			-	0.083	-	-		0.04				0.032		
Nutrients (mg/L)																				
Nitrate (NO3 as N)	0.7 (7)		0.05	0.11	0.13	0.38	0.02	0.16	0.03	0.06	0.05	0.02	0.01	0.15	0.03	0.05	0.04	0.22	0.06	<0.01
Nitrite (NO2 as N)			<0.01	0.12	0.02	0.01	<0.01	<0.01	<0.01	<0.01	0.02	0.15	0.1	0.04	0.24	<0.01	0.01	0.02	0.02	0.02
Ammonia (NH3 as N)	1.88 (2)	2.84 (2)	0.85	3.07	1.73	1.73	<0.10	<0.10	0.81	2.6	0.52	2.6	1.73	0.68	3.5	0.07	0.83	2.39	1.65	1.24
Total Kjeldahl Nitrogen (TKN)	0.5 (5)	0.12 (6)	1.2	4.1	2.5	1.9	0.2	1.7	1.7	3.1	0.6	2.7	2	1.2	4.3	1.6	1.7	2.6	2.7	2.5
Total Organic Carbon (TOC)									-	-	-	-								
Total Phosphorus (TP)	0.05 (7)	0.025 (7)	0.03	0.03	<0.01	0.02	<0.01	1.95	0.06	0.03	0.01	0.03	0.22	0.05	0.09	0.2	0.11	0.06	0.03	0.03
Biological (CFU/100 ml)																				
Enterococcus	35 (8)	35 (8)				12			-	~16		-		130				240		
Thermotolerant (Faecal) coliforms	150 (8)	150 (8)				4			-	~6	-	-		220				1100		

Exceeds ANZECC (2000) guidelines

0.054

Focus of this monitoring report


Table 6: Groundwater and Surface Water Monitoring - 2015 to 2016

		Groun	dwater			Blue Ang	gle Creek		Detection		
Analytes	27/05/15	24/08/15	4/11/15	26/02/16	27/05/15	24/08/15	4/11/15	26/02/16	Limit	Method Reference	
			Physica	l Propertie	es						
pН	X	X	X	X	X	X	X	X	0.01 pH unit	pH meter and probe/APHA4500-HB	
Electrical Conductivity	X	X	X	X	X	X	X	X	0.01 mS/cm	Conductivity meter and probe	
Dissolved Oxygen	X	X	X	X	X	X	X	X	0.0001	DO meter and probe	
Redox (Orp)		X				X			1 mV	Platinum electrode probe	
Temperature									1 °C	Temperature meter and probe	
Total Dissolved Solids	X	X	X	X	X	X	X	X	5 mg/L	Determined gravimetrically by drying (API 2540 C)	
Suspended Solids									2 mg/L	APHA2540D	
Turbidity									1 NTU	Turbidmeter	
			Nu	trients			•	•	•		
Ammonia-nitrogen	X	X	X	X	X	X	X	X	0.01 mg/L	FIA	
Total Phosphorus	X	X	X	X	X	X	X	X	2 μg/L	FIA	
Nitrate-nitrogen	X	X	X	X	X	X	X	X	10 μg/L	FIA	
Nitrite-nitrogen	X	X	X	X	X	X	X	X	1 μg/L	FIA	
Total Kjeldhal Nitrogen	X	X	X	X	X	X	X	X	50 μg/L	FIA	
			Hydro	-chemical							
Calcium		X				X			0.5 mg/L	USEPA 6010 A	
Chloride		X							0.5 mg/L	Titrated with mercuric nitrate using diphen carbazonel/xylene cyanol FF indicator	
Fluoride		X							0. 1 mg/L	APHA4500-FC	
Magnesium		X				X			0.02 mg/L	USEPA 6010 A	
Sulphate		X							1 mg/L	ICID/MS	
Sodium		X				X			0.05 mg/L	USEPA 6010 A	
Bicarbonate/Alkalinity	X	X	X	X	X	X	X	X	0.5 mg/L	APHA2340C	
Potassium		X				X			0.05 mg/L	USEPA 6010 A	
			Organic (Contamina	nts						
Dissolved Organic Carbon		X			I			I	0.50 mg/L	APHA 5310C	
Total Organic Carbon									0.1 mg/L	APHA 5310C	
C			Inorganic	Contamina	ants			<u> </u>			
Iron		X	-			X		<u> </u>	1 μg/L	USEPA 6010 A	
Manganese		X				X			1 μg/L	USEPA 6010	
			Biological	Contamin	ants						
Thermotolerant (Faecal) coliforms MF						X			1cfu/100 ml	WMM 009 (~AS 4276.7 - 1995)	
Enterococcus MF						X			1cfu/100 ml	WMM 013 (~AS 4276.9 - 1995)	




Graphs


Graph-01: Groundwater Ammonia Time-Series Trends - Shallow & Creek Wells




Graph -02: Groundwater Ammonia Time-Series Trends - Deep Wells




Graph-03: Groundwater Total Phosphorous Time-Series Trends - Standard Wells




Graph-04: Groundwater Total Phosphorous Time-Series Trends - New Shallow and Deep Wells





Graph-05: Depth to Groundwater (m AHD) - 2001 to February 2016





Sampling Date







Figures





Source: Neil Charters Pty Ltd

Date: Feb 2016


Reference: E2W-025_55.cdr

GERROA ANNUAL MONITORING REPORT (2015-2016)

Figure 1







KEY

MW1S Monitoring Well - Shallow, August 2006

MW1D Monitoring Well - Deep, August 2006

MW1

Standard Monitoring Well

ML1 • Surface Water Monitoring

Note: Bundled Piezometers MW1,6,7 - not sampled ML-3 no longer sampled, ML-2 and ML-4 no access




Source: URS Australia Pty Ltd

Reference: E2W-025_01.cdr

SITE LAYOUT & WELL LOCATIONS

Date: February 2016 GERROA ANNUAL MONITORING REPORT (2015-2016)






Source: URS Australia Pty Ltd- baseplan INFERRED GROUNDWATER FLOW REGIME (Wet, May 2015)

GERROA ANNUAL MONITORING REPORT (2015-2016)

Date: Feb 2016

Reference: E2W-025_59.cdr Figure 3A






Source: URS Australia Pty Ltd- baseplan INFERRED GROUNDWATER FLOW REGIME (Dry, August 2015)

GERROA ANNUAL MONITORING REPORT (2015-2016)

Date: February 2016
Reference: E2W-025_59.cdr



GERROA ANNUAL MONITORING REPORT (2015-2016)

Date: February 2016

Reference: E2W_025_58.cdr



Appendix A



CERTIFICATE OF ANALYSIS

Work Order : **EW1510398** Page : 1 of 6

Client : KIAMA COUNCIL Laboratory : Environmental Division NSW South Coast

Contact : MR PAUL CZULOWSKI Contact : Glenn Davies

Address : 11 MANNING STREET Address : 99 Kenny Street, Wollongong 2500

KIAMA NSW, AUSTRALIA 2533 Unit 4 / 13 Geary Place, PO Box 3105, North Nowra 2541

AUSTRALIA

E-mail : paulc@kiama.nsw.gov.au : glenn.davies@alsglobal.com

Telephone : +61 02 4232 0444 Telephone : 02 42253125

Facsimile : +61 02 4232 0555 Facsimile : W 02 42253128 N 02 44232083

Project : Gerroa Landfill Quarterly QC Level : NEPM 2013 Schedule B(3) and ALS QCS3 requirement

 Order number
 : --- Date Samples Received
 : 27-May-2015 17:30

 C-O-C number
 : --- Date Analysis Commenced
 : 27-May-2015

 Sampler
 : Craig Wilson
 Issue Date
 : 05-Jun-2015 16:18

Site · ----

No. of samples received : 17

Quote number : ---- No. of samples analysed : 17

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted.

This Certificate of Analysis contains the following information:

- General Comments
- Analytical Results



NATA Accredited Laboratory 825

Accredited for compliance with ISO/IEC 17025.

Signatories

This document has been electronically signed by the authorized signatories indicated below. Electronic signing has been carried out in compliance with procedures specified in 21 CFR Part 11.

Signatories Position Accreditation Category

Ankit Joshi Inorganic Chemist Sydney Inorganics
Glenn Davies Environmental Services Representative Laboratory - Wollongong

Page : 2 of 6
Work Order : EW1510398

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



General Comments

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the USEPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request.

Where moisture determination has been performed, results are reported on a dry weight basis.

Where a reported less than (<) result is higher than the LOR, this may be due to primary sample extract/digestate dilution and/or insufficient sample for analysis.

Where the LOR of a reported result differs from standard LOR, this may be due to high moisture content, insufficient sample (reduced weight employed) or matrix interference.

When sampling time information is not provided by the client, sampling dates are shown without a time component. In these instances, the time component has been assumed by the laboratory for processing purposes.

Key: CAS Number = CAS registry number from database maintained by Chemical Abstracts Services. The Chemical Abstracts Service is a division of the American Chemical Society.

LOR = Limit of reporting

^ = This result is computed from individual analyte detections at or above the level of reporting

ø = ALS is not NATA accredited for these tests.

- TDS by method EA-015 may bias high due to the presence of fine particulate matter, which may pass through the prescribed GF/C paper.
- Sampling and sample data supplied by ALS Wollongong.
- Sampling completed as per FWI-EN001 Groundwater Sampling.
- Sampling completed as per FWI-EN002 Surface Water Sampling.
- Field tests completed on day of sampling/receipt.

Page : 3 of 6
Work Order : EW1510398

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	MW1D	MW1S	MW3	MW4	MW5
	Cli	ent sampli	ng date / time	27-May-2015 14:45	27-May-2015 14:52	27-May-2015 14:37	27-May-2015 15:02	27-May-2015 14:31
Compound	CAS Number	LOR	Unit	EW1510398-001	EW1510398-002	EW1510398-003	EW1510398-004	EW1510398-005
				Result	Result	Result	Result	Result
EA005FD: Field pH								
pН		0.1	pH Unit	7.5	6.2	7.1	6.6	7.5
EA010FD: Field Conductivity								
Electrical Conductivity (Non		1	μS/cm	707	772	598	555	361
Compensated)								
EA015: Total Dissolved Solids								
Total Dissolved Solids @180°C		1	mg/L					
^ Total Dissolved Solids @180°C		1	mg/L	379	734	366	295	225
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1	<1	<1	<1	<1
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1	<1	<1	<1	<1
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	272	102	204	215	102
Total Alkalinity as CaCO3		1	mg/L	272	102	204	215	102
EK055G: Ammonia as N by Discrete	Analyser							
Ammonia as N	7664-41-7	0.01	mg/L	<0.01	0.09	0.15	0.75	0.03
EK057G: Nitrite as N by Discrete Ana	alyser							
Nitrite as N		0.01	mg/L	<0.01	<0.01	<0.01	<0.01	<0.01
EK058G: Nitrate as N by Discrete An	alyser							
^ Nitrate as N	14797-55-8	0.01	mg/L	1.87	<0.01	0.01	0.16	0.11
EK059G: Nitrite plus Nitrate as N (NC	Ox) by Discrete Ana	lyser						
Nitrite + Nitrate as N		0.01	mg/L	1.87	<0.01	0.01	0.16	0.11
EK061G: Total Kjeldahl Nitrogen By I	Discrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L	0.6	3.5	0.5	2.1	0.6
EK062G: Total Nitrogen as N (TKN +	NOx) by Discrete An	alvser						
^ Total Nitrogen as N		0.1	mg/L	2.5	3.5	0.5	2.3	0.7
EK067G: Total Phosphorus as P by D)iscrete Analyser							
Total Phosphorus as P		0.01	mg/L	0.05	0.26	0.10	1.44	0.09
EN67 PK: Field Tests								
Field Observations		0.01						
EP025FD: Field Dissolved Oxygen								
Dissolved Oxygen		0.01	mg/L	3,45	2.19	3.05	3.57	3.95
		0.01	my/L	Vrv	2.10	3.00	5.01	3.50
FWI-EN/001: Groundwater Sampling Depth	- Deptn 	0.01	m	2.92	3.06	3.39	3.70	3.26
Бериі		0.01	111	2.32	3.00	3.33	3.70	J.20

Page : 4 of 6
Work Order : EW1510398

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	MW6D	MW6S	MW7D	MW7S	MW9
	Cli	ent sampli	ng date / time	27-May-2015 14:05	27-May-2015 14:00	27-May-2015 14:16	27-May-2015 14:11	27-May-2015 15:25
Compound	CAS Number	LOR	Unit	EW1510398-006	EW1510398-007	EW1510398-008	EW1510398-009	EW1510398-010
				Result	Result	Result	Result	Result
EA005FD: Field pH								
pН		0.1	pH Unit	7.0	6.2	7.4	6.8	6.4
EA010FD: Field Conductivity								
Electrical Conductivity (Non		1	μS/cm	821	573	426	317	9120
Compensated)								
EA015: Total Dissolved Solids								
Total Dissolved Solids @180°C		1	mg/L				248	
^ Total Dissolved Solids @180°C		1	mg/L	468	320	249		3580
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1	<1	<1	<1	<1
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1	<1	<1	<1	<1
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	292	40	175	89	142
Total Alkalinity as CaCO3		1	mg/L	292	40	175	89	142
EK055G: Ammonia as N by Discrete	Analyser							
Ammonia as N	7664-41-7	0.01	mg/L	3.13	0.09	0.96	0.59	<0.01
EK057G: Nitrite as N by Discrete Ana	alyser							
Nitrite as N		0.01	mg/L	0.06	<0.01	<0.01	<0.01	<0.01
EK058G: Nitrate as N by Discrete An	alyser							
^ Nitrate as N	14797-55-8	0.01	mg/L	1.13	<0.01	0.46	0.04	<0.01
EK059G: Nitrite plus Nitrate as N (NC	Ox) by Discrete Ana	lyser						
Nitrite + Nitrate as N		0.01	mg/L	1.19	<0.01	0.46	0.04	<0.01
EK061G: Total Kjeldahl Nitrogen By I	Discrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L	3.2	2.9	1.3	0.8	2.4
EK062G: Total Nitrogen as N (TKN +	NOx) by Discrete An	alvser						
^ Total Nitrogen as N		0.1	mg/L	4.4	2.9	1.8	0.8	2.4
EK067G: Total Phosphorus as P by D)iscrete Analyser							
Total Phosphorus as P		0.01	mg/L	2.01	0.40	0.71	0.21	0.13
EN67 PK: Field Tests								
Field Observations		0.01						
EP025FD: Field Dissolved Oxygen								
Dissolved Oxygen		0.01	mg/L	2.38	3.48	3.40	3,22	3.61
		0.01	g, _	2.00	0.40	0.40	V.22	0.01
FWI-EN/001: Groundwater Sampling Depth	- Deptn 	0.01	m	4.47	4.20	4.21	4.06	1.66
Бериі		0.01	111	4.4/	4.20	4.21	4.00	1.00

Page : 5 of 6
Work Order : EW1510398

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	MW10	MW11	ML-1	ML-2	ML-3
	Cli	ent sampli	ng date / time	27-May-2015 15:15	27-May-2015 15:37	27-May-2015 15:05	27-May-2015 15:47	27-May-2015 15:07
Compound	CAS Number	LOR	Unit	EW1510398-011	EW1510398-012	EW1510398-013	EW1510398-014	EW1510398-015
				Result	Result	Result	Result	Result
EA005FD: Field pH								
pH		0.1	pH Unit	5.5	5.9		6.6	
EA010FD: Field Conductivity								
Electrical Conductivity (Non		1	μS/cm	8540	4200		4280	
Compensated)								
EA015: Total Dissolved Solids								
Total Dissolved Solids @180°C		1	mg/L					
^ Total Dissolved Solids @180°C		1	mg/L	3530	2290		2060	
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1	<1		<1	
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1	<1		<1	
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	16	68		68	
Total Alkalinity as CaCO3		1	mg/L	16	68		68	
EK055G: Ammonia as N by Discrete	Analyser							
Ammonia as N	7664-41-7	0.01	mg/L	0.05	<0.01		0.80	
EK057G: Nitrite as N by Discrete Ana	alyser							
Nitrite as N		0.01	mg/L	<0.01	<0.01		0.02	
EK058G: Nitrate as N by Discrete An	alyser							
^ Nitrate as N	14797-55-8	0.01	mg/L	<0.01	<0.01		0.02	
EK059G: Nitrite plus Nitrate as N (NC	Dx) by Discrete Ana	lyser						
Nitrite + Nitrate as N		0.01	mg/L	<0.01	<0.01		0.04	
EK061G: Total Kjeldahl Nitrogen By I	Discrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L	3.2	5.6		1.2	
EK062G: Total Nitrogen as N (TKN +	NOx) by Discrete An	alvser						
^ Total Nitrogen as N		0.1	mg/L	3.2	5.6		1.2	
EK067G: Total Phosphorus as P by D)iscrete Analyser							
Total Phosphorus as P		0.01	mg/L	0.26	0.16		0.16	
EN67 PK: Field Tests								
Field Observations		0.01				No Access		No Access
EP025FD: Field Dissolved Oxygen								
Dissolved Oxygen		0.01	mg/L	3.29	2.11		4.46	
FWI-EN/001: Groundwater Sampling			J. =					
Depth	- Deptn 	0.01	m	2.07	2.08			
Sopui		0.01		2.01	2.00			

Page : 6 of 6
Work Order : EW1510398

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Cli	ent sample ID	ML-4	ML-5			
	Clie	ent sampli	ng date / time	27-May-2015 15:09	27-May-2015 15:30			
Compound	CAS Number	LOR	Unit	EW1510398-016	EW1510398-017			
				Result	Result	Result	Result	Result
EA005FD: Field pH								
рН		0.1	pH Unit		6.9			
EA010FD: Field Conductivity								
Electrical Conductivity (Non		1	μS/cm		1210			
Compensated)								
EA015: Total Dissolved Solids								
Total Dissolved Solids @180°C		1	mg/L					
^ Total Dissolved Solids @180°C		1	mg/L		538			
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L		<1			
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L		<1			
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L		60			
Total Alkalinity as CaCO3		1	mg/L		60			
EK055G: Ammonia as N by Discrete								
Ammonia as N	7664-41-7	0.01	mg/L		0.83			
EK057G: Nitrite as N by Discrete Ana	alyser							
Nitrite as N		0.01	mg/L		0.01			
EK058G: Nitrate as N by Discrete An	alyser							
^ Nitrate as N	14797-55-8	0.01	mg/L		0.04			
EK059G: Nitrite plus Nitrate as N (NC	Dx) by Discrete Anal	yser						
Nitrite + Nitrate as N		0.01	mg/L		0.05			
EK061G: Total Kjeldahl Nitrogen By I	Discrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L		1.7			
EK062G: Total Nitrogen as N (TKN +	NOx) by Discrete An	alyser						
^ Total Nitrogen as N		0.1	mg/L		1.8			
EK067G: Total Phosphorus as P by D	Discrete Analyser							
Total Phosphorus as P		0.01	mg/L		0.11			
EN67 PK: Field Tests								
Field Observations		0.01		No Access				
EP025FD: Field Dissolved Oxygen								
Dissolved Oxygen		0.01	mg/L		5.20			
FWI-EN/001: Groundwater Sampling								
Depth	- Deptii	0.01	m					
P						1		



CERTIFICATE OF ANALYSIS

Work Order Page : EW1511400 : 1 of 11

Client : KIAMA COUNCIL Laboratory : Environmental Division NSW South Coast

: MR PAUL CZULOWSKI Contact Contact : Glenn Davies

Address Address : 11 MANNING STREET : 1/19 Ralph Black Drive, KIAMA NSW, AUSTRALIA 2533

North Wollongong, 2500

E-mail E-mail : paulc@kiama.nsw.gov.au : glenn.davies@alsglobal.com

: +61 02 4232 0444 : 02 42253125 Telephone Telephone

: +61 02 4232 0555 Facsimile : W 02 42253128 N 02 44232083 Facsimile

Project : Gerroa Landfill Annual QC Level : NEPM 2013 Schedule B(3) and ALS QCS3 requirement

Order number : 87895 Date Samples Received : 24-Aug-2015 14:40 C-O-C number Date Analysis Commenced

: 24-Aug-2015 Sampler Issue Date : Craig Wilson : 01-Sep-2015 16:54

Site

No. of samples received : 18 Quote number No. of samples analysed · 18

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted.

This Certificate of Analysis contains the following information:

- General Comments
- Analytical Results

Page Work Order Client

Project

: 2 of 11 EW1511400 : KIAMA COUNCIL

Gerroa Landfill Annual

WORLD RECOGNISED **ACCREDITATION**

NATA Accredited Laboratory 825

Accredited for compliance with ISO/IEC 17025.

Signatories

This document has been electronically signed by the authorized signatories indicated below. Electronic signing has been carried out in compliance with procedures specified in 21 CFR Part 11.

Signatories	Position	Accreditation Category
Ankit Joshi	Inorganic Chemist	Sydney Inorganics
Ashesh Patel	Inorganic Chemist	Sydney Inorganics
Celine Conceicao	Senior Spectroscopist	Sydney Inorganics
Glenn Davies	Environmental Services Representative	Laboratory - Wollongong
Raymond Commodore	Instrument Chemist	Sydney Inorganics
Somlok Chai	Microbiologist	Sydney Microbiology

Page : 3 of 11 Work Order : EW1511400

Client : KIAMA COUNCIL
Project : Gerroa Landfill Annual



General Comments

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the USEPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request.

Where moisture determination has been performed, results are reported on a dry weight basis.

Where a reported less than (<) result is higher than the LOR, this may be due to primary sample extract/digestate dilution and/or insufficient sample for analysis.

Where the LOR of a reported result differs from standard LOR, this may be due to high moisture content, insufficient sample (reduced weight employed) or matrix interference.

When sampling time information is not provided by the client, sampling dates are shown without a time component. In these instances, the time component has been assumed by the laboratory for processing purposes.

Key: CAS Number = CAS registry number from database maintained by Chemical Abstracts Services. The Chemical Abstracts Service is a division of the American Chemical Society.

LOR = Limit of reporting

^ = This result is computed from individual analyte detections at or above the level of reporting

ø = ALS is not NATA accredited for these tests.

- TDS by method EA-015 may bias high due to the presence of fine particulate matter, which may pass through the prescribed GF/C paper.
- Ionic Balance out of acceptable limits due to analytes not quantified in this report.
- Sampling and sample data supplied by ALS Wollongong.
- Sampling completed as per FWI-EN001 Groundwater Sampling.
- Field tests completed on day of sampling/receipt.
- MW023 is ALS's internal code and is equivalent to AS4276.9.
- MW006 is ALS's internal code and is equivalent to AS4276.7.

Page : 4 of 11 Work Order : EW1511400

Client : KIAMA COUNCIL
Project : Gerroa Landfill Annual



Sub-Matrix: WATER (Matrix: WATER)		Cli	ent sample ID	MW1D	MW1S	MW3	MW4	MW5
	CI	lient sampli	ing date / time	24-Aug-2015 12:00	24-Aug-2015 12:10	24-Aug-2015 11:30	24-Aug-2015 09:15	24-Aug-2015 11:05
Compound	CAS Number	LOR	Unit	EW1511400-001	EW1511400-002	EW1511400-003	EW1511400-004	EW1511400-005
·				Result	Result	Result	Result	Result
EA005FD: Field pH								
pH		0.1	pH Unit	7.7		7.3	7.0	8.0
EA010FD: Field Conductivity								
Electrical Conductivity (Non Compensated)		1	μS/cm	709		962	555	331
EA015: Total Dissolved Solids								
^ Total Dissolved Solids @180°C		1	mg/L	443		565	330	199
EA075FD: Field Redox Potential								
Redox Potential		0.1	mV	-37.0		-138	76.0	56.0
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1		<1	<1	<1
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1		<1	<1	<1
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	262		250	230	118
Total Alkalinity as CaCO3		1	mg/L	262		250	230	118
ED041G: Sulfate (Turbidimetric) as SO4	1 2- by DA							
Sulfate as SO4 - Turbidimetric	14808-79-8	1	mg/L	10		13	9	5
ED045G: Chloride by Discrete Analyser								
Chloride	16887-00-6	1	mg/L	37		106	18	18
ED093F: Dissolved Major Cations								
Calcium	7440-70-2	1	mg/L	72		98	90	44
Magnesium	7439-95-4	1	mg/L	24		13	6	5
Sodium	7440-23-5	1	mg/L	37		73	14	12
Potassium	7440-09-7	1	mg/L	13		3	2	2
ED093T: Total Major Cations								
Calcium	7440-70-2	1	mg/L					
Magnesium	7439-95-4	1	mg/L					
Sodium	7440-23-5	1	mg/L					
Potassium	7440-09-7	1	mg/L					
EG020F: Dissolved Metals by ICP-MS								
Manganese	7439-96-5	0.001	mg/L	0.006		0.184	0.086	<0.001
Iron	7439-89-6	0.05	mg/L	<0.05		15.5	<0.05	<0.05
EG020T: Total Metals by ICP-MS						·		
Manganese	7439-96-5	0.001	mg/L					
Iron	7439-89-6	0.05	mg/L					

Page : 5 of 11 Work Order : EW1511400

Client : KIAMA COUNCIL
Project : Gerroa Landfill Annual



Sub-Matrix: WATER (Matrix: WATER)		CI	ient sample ID	MW1D	MW1S	MW3	MW4	MW5
	Clie	nt sampl	ling date / time	24-Aug-2015 12:00	24-Aug-2015 12:10	24-Aug-2015 11:30	24-Aug-2015 09:15	24-Aug-2015 11:05
Compound	CAS Number	LOR	Unit	EW1511400-001	EW1511400-002	EW1511400-003	EW1511400-004	EW1511400-005
·				Result	Result	Result	Result	Result
EK040P: Fluoride by PC Titrator								
Fluoride	16984-48-8	0.1	mg/L	0.2		0.1	0.2	0.2
EK055G: Ammonia as N by Discrete A	Analyser							
Ammonia as N	7664-41-7	0.01	mg/L	0.77		0.28	0.15	0.02
EK057G: Nitrite as N by Discrete Ana	ılyser							
Nitrite as N	14797-65-0	0.01	mg/L	0.04		<0.01	<0.01	<0.01
EK058G: Nitrate as N by Discrete Ana	alyser							
^ Nitrate as N	14797-55-8	0.01	mg/L	2.48		0.02	0.03	0.09
EK059G: Nitrite plus Nitrate as N (NO	x) by Discrete Analy	/ser						
Nitrite + Nitrate as N		0.01	mg/L	2.52		0.02	0.03	0.09
EK061G: Total Kjeldahl Nitrogen By D	Discrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L	1.1		0.7	0.6	0.3
EK067G: Total Phosphorus as P by D	iscrete Analyser							
Total Phosphorus as P		0.01	mg/L	0.06		0.46	2.55	0.06
EN055: Ionic Balance								
Total Anions		0.01	meq/L	6.62				
^ Total Anions		0.01	meq/L			8.26	5.29	2.97
^ Total Cations		0.01	meq/L	7.51		9.21	5.64	3.18
Ionic Balance		0.01	%	6.62				
^ Ionic Balance		0.01	%			5.49	3.27	3.45
EN67 PK: Field Tests								
Field Observations		0.01			DRY			
EP002: Dissolved Organic Carbon (Do	OC)							
Dissolved Organic Carbon		1	mg/L	9		12	3	4
EP025FD: Field Dissolved Oxygen								
Dissolved Oxygen		0.01	mg/L	1.69		3.05	3.21	3.39
FWI-EN/001: Groundwater Sampling -	Depth							
Depth		0.01	m	3.36		3.84	4.16	3.70
MW006: Faecal Coliforms & E.coli by	MF							
Faecal Coliforms		1	CFU/100mL					
MW023: Enterococci by Membrane Fi	Itration							
Enterococci		1	CFU/100mL					

Page : 6 of 11 Work Order : EW1511400

Client : KIAMA COUNCIL
Project : Gerroa Landfill Annual



Sub-Matrix: WATER (Matrix: WATER)		Cli	ent sample ID	MW6D	MW6S	MW7D	MW7S	MW9
	CI	ient sampli	ing date / time	24-Aug-2015 09:50	24-Aug-2015 09:30	24-Aug-2015 10:45	24-Aug-2015 10:20	24-Aug-2015 12:50
Compound	CAS Number	LOR	Unit	EW1511400-006	EW1511400-007	EW1511400-008	EW1511400-009	EW1511400-010
				Result	Result	Result	Result	Result
EA005FD: Field pH								
pH		0.1	pH Unit	7.2		7.4	8.1	6.7
EA010FD: Field Conductivity								
Electrical Conductivity (Non Compensated)		1	μS/cm	801		483	463	7690
EA015: Total Dissolved Solids								
^ Total Dissolved Solids @180°C		1	mg/L	522		300	267	5160
EA075FD: Field Redox Potential								
Redox Potential		0.1	mV	30.0		66.0	-75.0	130
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1		<1	<1	<1
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1		<1	<1	<1
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	305		214	140	135
Total Alkalinity as CaCO3		1	mg/L	305		214	140	135
ED041G: Sulfate (Turbidimetric) as SO4	2- by DA							
Sulfate as SO4 - Turbidimetric	14808-79-8	1	mg/L	69		7	12	384
ED045G: Chloride by Discrete Analyser								
Chloride	16887-00-6	1	mg/L	17		9	34	1960
ED093F: Dissolved Major Cations								
Calcium	7440-70-2	1	mg/L	139		75	42	69
Magnesium	7439-95-4	1	mg/L	11		7	6	165
Sodium	7440-23-5	1	mg/L	13		9	41	1350
Potassium	7440-09-7	1	mg/L	6		4	3	60
ED093T: Total Major Cations								
Calcium	7440-70-2	1	mg/L					
Magnesium	7439-95-4	1	mg/L					
Sodium	7440-23-5	1	mg/L					
Potassium	7440-09-7	1	mg/L					
EG020F: Dissolved Metals by ICP-MS								
Manganese	7439-96-5	0.001	mg/L	0.082		0.037	0.010	0.004
Iron	7439-89-6	0.05	mg/L	0.07		<0.05	0.42	0.26
EG020T: Total Metals by ICP-MS								
Manganese	7439-96-5	0.001	mg/L					
Iron	7439-89-6	0.05	mg/L					

Page : 7 of 11 Work Order : EW1511400

Client : KIAMA COUNCIL
Project : Gerroa Landfill Annual



Sub-Matrix: WATER (Matrix: WATER)		CI	ient sample ID	MW6D	MW6S	MW7D	MW7S	MW9
	Clie	nt sampl	ling date / time	24-Aug-2015 09:50	24-Aug-2015 09:30	24-Aug-2015 10:45	24-Aug-2015 10:20	24-Aug-2015 12:50
Compound	CAS Number	LOR	Unit	EW1511400-006	EW1511400-007	EW1511400-008	EW1511400-009	EW1511400-010
·				Result	Result	Result	Result	Result
EK040P: Fluoride by PC Titrator								
Fluoride	16984-48-8	0.1	mg/L	0.4		0.2	0.1	0.3
EK055G: Ammonia as N by Discrete A	Analyser							
Ammonia as N	7664-41-7	0.01	mg/L	1.77		1.38	0.54	0.02
EK057G: Nitrite as N by Discrete Ana	llyser							
Nitrite as N	14797-65-0	0.01	mg/L	0.05		0.01	<0.01	<0.01
EK058G: Nitrate as N by Discrete Ana	alyser							
^ Nitrate as N	14797-55-8	0.01	mg/L	1.40		0.33	<0.01	0.05
EK059G: Nitrite plus Nitrate as N (NC	x) by Discrete Analy	/ser						
Nitrite + Nitrate as N		0.01	mg/L	1.45		0.34	<0.01	0.05
EK061G: Total Kjeldahl Nitrogen By D	Discrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L	2.0		2.2	0.8	0.8
EK067G: Total Phosphorus as P by D	iscrete Analyser							
Total Phosphorus as P		0.01	mg/L	1.39		1.81	0.19	0.08
EN055: Ionic Balance								
Total Anions		0.01	meq/L					
^ Total Anions		0.01	meq/L	8.01		4.68	4.01	66.0
^ Total Cations		0.01	meq/L	8.56		4.81	4.45	77.3
Ionic Balance		0.01	%					
^ Ionic Balance		0.01	%	3.36		1.48	5.25	7.87
EN67 PK: Field Tests								
Field Observations		0.01			DRY			
EP002: Dissolved Organic Carbon (Do	OC)							
Dissolved Organic Carbon		1	mg/L	8		4	5	29
EP025FD: Field Dissolved Oxygen								
Dissolved Oxygen		0.01	mg/L	2.02		2.51	2.06	3.24
FWI-EN/001: Groundwater Sampling -	Depth							
Depth		0.01	m	4.88		4.63	4.50	1.90
MW006: Faecal Coliforms & E.coli by	MF							
Faecal Coliforms		1	CFU/100mL					
MW023: Enterococci by Membrane Fi	Itration							
Enterococci		1	CFU/100mL					

Page : 8 of 11 Work Order : EW1511400

Client : KIAMA COUNCIL
Project : Gerroa Landfill Annual



Sub-Matrix: WATER (Matrix: WATER)		Cli	ent sample ID	MW10	MW11	ML-1	ML-2	ML-3
	Cli	ient sampli	ing date / time	24-Aug-2015 12:25	24-Aug-2015 13:25	24-Aug-2015 13:50	24-Aug-2015 13:35	24-Aug-2015 13:55
Compound	CAS Number	LOR	Unit	EW1511400-011	EW1511400-012	EW1511400-013	EW1511400-014	EW1511400-015
				Result	Result	Result	Result	Result
EA005FD: Field pH								
рН		0.1	pH Unit		6.4		7.5	
EA010FD: Field Conductivity								
Electrical Conductivity (Non Compensated)		1	μS/cm		1730		7790	
EA015: Total Dissolved Solids								
^ Total Dissolved Solids @180°C		1	mg/L		1470		4840	
EA075FD: Field Redox Potential								
Redox Potential		0.1	mV		-30.0		-17.0	
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L		<1		<1	
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L		<1		<1	
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L		83		103	
Total Alkalinity as CaCO3		1	mg/L		83		103	
ED041G: Sulfate (Turbidimetric) as SO4	1 2- by DA							
Sulfate as SO4 - Turbidimetric	14808-79-8	1	mg/L		41			
ED045G: Chloride by Discrete Analyser								
Chloride	16887-00-6	1	mg/L		374			
ED093F: Dissolved Major Cations								
Calcium	7440-70-2	1	mg/L		10			
Magnesium	7439-95-4	1	mg/L		23			
Sodium	7440-23-5	1	mg/L		302			
Potassium	7440-09-7	1	mg/L		13			
ED093T: Total Major Cations								
Calcium	7440-70-2	1	mg/L				66	
Magnesium	7439-95-4	1	mg/L				158	
Sodium	7440-23-5	1	mg/L				1420	
Potassium	7440-09-7	1	mg/L				55	
EG020F: Dissolved Metals by ICP-MS								
Manganese	7439-96-5	0.001	mg/L		0.004			
Iron	7439-89-6	0.05	mg/L		1.28			
EG020T: Total Metals by ICP-MS								
Manganese	7439-96-5	0.001	mg/L				0.023	
Iron	7439-89-6	0.05	mg/L				1.64	

Page : 9 of 11 Work Order : EW1511400

Client : KIAMA COUNCIL
Project : Gerroa Landfill Annual



Sub-Matrix: WATER (Matrix: WATER)		Cli	ient sample ID	MW10	MW11	ML-1	ML-2	ML-3
	Clie	nt sampl	ing date / time	24-Aug-2015 12:25	24-Aug-2015 13:25	24-Aug-2015 13:50	24-Aug-2015 13:35	24-Aug-2015 13:55
Compound	CAS Number	LOR	Unit	EW1511400-011	EW1511400-012	EW1511400-013	EW1511400-014	EW1511400-015
				Result	Result	Result	Result	Result
EK040P: Fluoride by PC Titrator								
Fluoride	16984-48-8	0.1	mg/L		0.3			
EK055G: Ammonia as N by Discrete A	nalyser							
Ammonia as N	7664-41-7	0.01	mg/L		0.01		1.50	
EK057G: Nitrite as N by Discrete Ana	lyser							
Nitrite as N	14797-65-0	0.01	mg/L		<0.01		0.02	
EK058G: Nitrate as N by Discrete Ana	llyser							
^ Nitrate as N	14797-55-8	0.01	mg/L		<0.01		0.13	
EK059G: Nitrite plus Nitrate as N (NO	x) by Discrete Analy	/ser						
Nitrite + Nitrate as N		0.01	mg/L		<0.01		0.15	
EK061G: Total Kjeldahl Nitrogen By D	iscrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L		4.9		1.9	
EK067G: Total Phosphorus as P by Di	screte Analyser							
Total Phosphorus as P		0.01	mg/L		0.31		0.10	
EN055: Ionic Balance								
Total Anions		0.01	meq/L					
^ Total Anions		0.01	meq/L		13.1			
^ Total Cations		0.01	meq/L		15.9			
Ionic Balance		0.01	%					
^ Ionic Balance		0.01	%		9.66			
EN67 PK: Field Tests								
Field Observations		0.01		DRY		NO ACCESS		NO ACCESS
EP002: Dissolved Organic Carbon (DO	OC)							
Dissolved Organic Carbon		1	mg/L		102			
EP025FD: Field Dissolved Oxygen								
Dissolved Oxygen		0.01	mg/L		1.80		9.11	
FWI-EN/001: Groundwater Sampling -	Depth							
Depth		0.01	m		2.33			
MW006: Faecal Coliforms & E.coli by	MF							
Faecal Coliforms		1	CFU/100mL				640	
MW023: Enterococci by Membrane Fil	tration							
Enterococci		1	CFU/100mL				590	

Page : 10 of 11 Work Order : EW1511400

Client : KIAMA COUNCIL
Project : Gerroa Landfill Annual



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	ML-4	ML-5	BLANK		
	CI	ient sampli	ng date / time	24-Aug-2015 13:40	24-Aug-2015 13:05	24-Aug-2015 10:00		
Compound	CAS Number	LOR	Unit	EW1511400-016	EW1511400-017	EW1511400-018		
				Result	Result	Result	Result	Result
EA005FD: Field pH								
рН		0.1	pH Unit		7.3			
EA010FD: Field Conductivity								
Electrical Conductivity (Non Compensated)		1	μS/cm		2380			
EA015: Total Dissolved Solids								
^ Total Dissolved Solids @180°C		1	mg/L		1370			
EA075FD: Field Redox Potential								
Redox Potential		0.1	mV		95.0			
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L		<1			
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L		<1			
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L		134			
Total Alkalinity as CaCO3		1	mg/L		134			
ED041G: Sulfate (Turbidimetric) as SO4	2- by DA							
Sulfate as SO4 - Turbidimetric	14808-79-8	1	mg/L					
ED045G: Chloride by Discrete Analyser								
Chloride	16887-00-6	1	mg/L					
ED093F: Dissolved Major Cations								
Calcium	7440-70-2	1	mg/L			<1		
Magnesium	7439-95-4	1	mg/L			<1		
Sodium	7440-23-5	1	mg/L			<1		
Potassium	7440-09-7	1	mg/L			<1		
ED093T: Total Major Cations								
Calcium	7440-70-2	1	mg/L		42			
Magnesium	7439-95-4	1	mg/L		49			
Sodium	7440-23-5	1	mg/L		379			
Potassium	7440-09-7	1	mg/L		19			
EG020F: Dissolved Metals by ICP-MS								
Manganese	7439-96-5	0.001	mg/L			<0.001		
Iron	7439-89-6	0.05	mg/L			<0.05		
EG020T: Total Metals by ICP-MS								
Manganese	7439-96-5	0.001	mg/L		0.032			
Iron	7439-89-6	0.05	mg/L		2.09			

Page : 11 of 11 Work Order : EW1511400

Client : KIAMA COUNCIL
Project : Gerroa Landfill Annual



Compound CaS Number LOR Unit EWIST1400-015 EWIST14	Sub-Matrix: WATER (Matrix: WATER)		Cli	ient sample ID	ML-4	ML-5	BLANK		
Result R		Clie	nt sampl	ing date / time	24-Aug-2015 13:40	24-Aug-2015 13:05	24-Aug-2015 10:00		
EK040P: Fluoride by PC Titrator Fluoride Fluoride	Compound	CAS Number	LOR	Unit	EW1511400-016	EW1511400-017	EW1511400-018		
Fluoride					Result	Result	Result	Result	Result
Fluoride	EK040P: Fluoride by PC Titrator								
Ammonia as N		16984-48-8	0.1	mg/L					
Ammonia as N	EK055G: Ammonia as N by Discrete An	alyser							
Nitrite as N 14797-65-0 0.01 mg/L 0.02			0.01	mg/L		2.39			
Nitrite as N	EK057G: Nitrite as N by Discrete Analys	ser							
^ Nitrate as N			0.01	mg/L		0.02			
^ Nitrate as N	EK058G: Nitrate as N by Discrete Analy	/ser							
Nitrite + Nitrate as N			0.01	mg/L		0.22			
Nitrite + Nitrate as N	EK059G: Nitrite plus Nitrate as N (NOx)	by Discrete Analy	/ser						
Total Kjeldahi Nitrogen as N				mg/L		0.24			
Total Kjeldahl Nitrogen as N	EK061G: Total Kieldahl Nitrogen By Dis	crete Analyser							
Total Phosphorus as P			0.1	mg/L		2.6			
Total Phosphorus as P	EK067G: Total Phosphorus as P by Disc	crete Analyser							
Total Anions			0.01	mg/L		0.06			
Total Anions	EN055: Ionic Balance								
^ Total Cations 0.01 meq/L			0.01	meq/L					
Ionic Balance	^ Total Anions		0.01	meq/L					
^ Ionic Balance	^ Total Cations		0.01	meq/L					
EN67 PK: Field Tests Field Observations 0.01 NO ACCESS EP002: Dissolved Organic Carbon (DOC) Dissolved Organic Carbon 1 mg/L	Ionic Balance		0.01	%					
Field Observations	^ Ionic Balance		0.01	%					
EP002: Dissolved Organic Carbon (DOC) Dissolved Organic Carbon 1 mg/L	EN67 PK: Field Tests								
Dissolved Organic Carbon	Field Observations		0.01		NO ACCESS				
EP025FD: Field Dissolved Oxygen	EP002: Dissolved Organic Carbon (DOC	;)							
Dissolved Oxygen	Dissolved Organic Carbon		1	mg/L					
Dissolved Oxygen	EP025FD: Field Dissolved Oxygen								
Depth 0.01 m -			0.01	mg/L		7.34			
Depth 0.01 m -	FWI-EN/001: Groundwater Sampling - D	epth							
Faecal Coliforms 1 CFU/100mL 1100 MW023: Enterococci by Membrane Filtration			0.01	m					
Faecal Coliforms 1 CFU/100mL 1100 MW023: Enterococci by Membrane Filtration	MW006: Faecal Coliforms & E.coli by M								
			1	CFU/100mL		1100			
	MW023: Enterococci by Membrane Filtr	ation							
Enterococci 1 CFU/100mL 240			1	CFU/100mL		240			



CERTIFICATE OF ANALYSIS

Work Order : **EW1512393** Page : 1 of 6

Client : KIAMA COUNCIL Laboratory : Environmental Division NSW South Coast

Contact : MR PAUL CZULOWSKI Contact : Glenn Davies

Address : 11 MANNING STREET Address : 1/19 Ralph Black Dr, North Wollongong 2500

4/13 Geary PI, North Nowra 2541

Australia

Telephone : +61 02 4232 0444 Telephone : 02 42253125

Facsimile : +61 02 4232 0555 Facsimile : W 02 42253128 N 02 44232083

Project : Gerroa Landfill Quarterly QC Level : NEPM 2013 Schedule B(3) and ALS QCS3 requirement

 Order number
 : 87895
 Date Samples Received
 : 04-Nov-2015 10:13

 C-O-C number
 : --- Date Analysis Commenced
 : 04-Nov-2015

Sampler : Craig Wilson | Issue Date : 11-Nov-2015 11:53

Site · ----

Quote number : ---- No. of samples received : 17

Quote number : ---- No. of samples analysed : 17

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted.

KIAMA NSW, AUSTRALIA 2533

This Certificate of Analysis contains the following information:

- General Comments
- Analytical Results



NATA Accredited Laboratory 825

Accredited for compliance with ISO/IEC 17025.

Signatories

This document has been electronically signed by the authorized signatories indicated below. Electronic signing has been carried out in compliance with procedures specified in 21 CFR Part 11.

Signatories Position Accreditation Category

Ankit Joshi Inorganic Chemist Sydney Inorganics
Dian Dao Sydney Inorganics
Glenn Davies Environmental Services Representative Laboratory - Wollongong

Page : 2 of 6
Work Order : EW1512393

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



General Comments

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the USEPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request.

Where moisture determination has been performed, results are reported on a dry weight basis.

Where a reported less than (<) result is higher than the LOR, this may be due to primary sample extract/digestate dilution and/or insufficient sample for analysis.

Where the LOR of a reported result differs from standard LOR, this may be due to high moisture content, insufficient sample (reduced weight employed) or matrix interference.

When sampling time information is not provided by the client, sampling dates are shown without a time component. In these instances, the time component has been assumed by the laboratory for processing purposes.

Key: CAS Number = CAS registry number from database maintained by Chemical Abstracts Services. The Chemical Abstracts Service is a division of the American Chemical Society.

LOR = Limit of reporting

^ = This result is computed from individual analyte detections at or above the level of reporting

ø = ALS is not NATA accredited for these tests.

- TDS by method EA-015 may bias high for sample 2 due to the presence of fine particulate matter, which may pass through the prescribed GF/C paper.
- Sampling and sample data supplied by ALS Wollongong.
- Sampling completed as per FWI-EN001 Groundwater Sampling.
- Sampling completed as per FWI-EN002 Surface Water Sampling.
- Field tests completed on day of sampling/receipt.

Page : 3 of 6
Work Order : EW1512393

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	MW1D	MW1S	MW3	MW4	MW5
	Cli	ent sampli	ng date / time	04-Nov-2015 10:20	04-Nov-2015 10:35	04-Nov-2015 10:10	04-Nov-2015 10:55	04-Nov-2015 10:00
Compound	CAS Number	LOR	Unit	EW1512393-001	EW1512393-002	EW1512393-003	EW1512393-004	EW1512393-005
				Result	Result	Result	Result	Result
EA005FD: Field pH								
рН		0.1	pH Unit	7.3	6.0	7.2	6.4	7.8
EA010FD: Field Conductivity								
Electrical Conductivity (Non		1	μS/cm	803	342	843	418	410
Compensated)								
EA015: Total Dissolved Solids								
Total Dissolved Solids @180°C		1	mg/L	480	385	488	258	254
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1	<1	<1	<1	<1
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1	<1	<1	<1	<1
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	291	51	169	178	147
Total Alkalinity as CaCO3		1	mg/L	291	51	169	178	147
EK055G: Ammonia as N by Discrete	Analyser							
Ammonia as N	7664-41-7	0.01	mg/L	3.78	0.23	0.74	0.26	0.03
EK057G: Nitrite as N by Discrete Ana	alyser							
Nitrite as N	14797-65-0	0.01	mg/L	<0.01	<0.01	0.01	<0.01	0.06
EK058G: Nitrate as N by Discrete An	alvser							
Nitrate as N	14797-55-8	0.01	mg/L	0.13	<0.01	0.02	<0.01	0.07
EK059G: Nitrite plus Nitrate as N (NC	Dx) by Discrete Anal	lvser						
Nitrite + Nitrate as N		0.01	mg/L	0.13	<0.01	0.03	<0.01	0.13
EK061G: Total Kjeldahl Nitrogen By [Discrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L	4.6	2.1	2.1	0.7	0.4
EK062G: Total Nitrogen as N (TKN +	NOx) by Discrete An	alvser						
^ Total Nitrogen as N		0.1	mg/L	4.7	2.1	2.1	0.7	0.5
EK067G: Total Phosphorus as P by D	Discrete Analyser		3				-	
Total Phosphorus as P		0.01	mg/L	0.06	0.18	0.83	1.97	0.02
EN67 PK: Field Tests			J. –					,
Field Observations		0.01						
		0.01						<u> </u>
EP025FD: Field Dissolved Oxygen Dissolved Oxygen		0.01	mg/L	1.71	2.20	1.54	3.03	2.03
		0.01	my/L	1./ 1	2.20	1.04	3.03	2.03
FWI-EN/001: Groundwater Sampling		0.01	m	3.09	3.20	3.55	3.90	3.46
Depth		0.01	m	3.09	3.20	3.55	3.90	3.46

Page : 4 of 6
Work Order : EW1512393

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	MW6D	MW6S	MW7D	MW7S	MW9
	Clie	ent sampli	ng date / time	04-Nov-2015 09:25	04-Nov-2015 09:20	04-Nov-2015 09:35	04-Nov-2015 09:45	04-Nov-2015 11:20
Compound	CAS Number	LOR	Unit	EW1512393-006	EW1512393-007	EW1512393-008	EW1512393-009	EW1512393-010
				Result	Result	Result	Result	Result
EA005FD: Field pH								
рН		0.1	pH Unit	7.0		7.3	7.1	6.5
EA010FD: Field Conductivity								
Electrical Conductivity (Non		1	μS/cm	953		499	423	4610
Compensated)								
EA015: Total Dissolved Solids								
Total Dissolved Solids @180°C		1	mg/L	574		337	357	2870
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1		<1	<1	<1
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1		<1	<1	<1
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	347		215	133	133
Total Alkalinity as CaCO3		1	mg/L	347		215	133	133
EK055G: Ammonia as N by Discrete	Analyser							
Ammonia as N	7664-41-7	0.01	mg/L	4.53		1.64	0.80	0.47
EK057G: Nitrite as N by Discrete Ana	alyser							
Nitrite as N	14797-65-0	0.01	mg/L	<0.01		0.03	<0.01	<0.01
EK058G: Nitrate as N by Discrete An	alvser							
Nitrate as N	14797-55-8	0.01	mg/L	<0.01		<0.01	<0.01	<0.01
EK059G: Nitrite plus Nitrate as N (NC	Dx) by Discrete Anal	vser						
Nitrite + Nitrate as N		0.01	mg/L	<0.01		0.03	<0.01	<0.01
EK061G: Total Kjeldahl Nitrogen By [Discrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L	6.0		2.3	1.3	2.2
EK062G: Total Nitrogen as N (TKN +	NOx) by Discrete An	alvser	-					
^ Total Nitrogen as N		0.1	mg/L	6.0		2.3	1.3	2.2
EK067G: Total Phosphorus as P by D	Discrete Analyser		3					
Total Phosphorus as P		0.01	mg/L	12.0		8.14	0.16	0.16
EN67 PK: Field Tests						2		
Field Observations		0.01			DRY			
		0.01						
EP025FD: Field Dissolved Oxygen Dissolved Oxygen		0.01	mg/L	1.47		1.95	1.82	1.92
		0.01	mg/L	1.47		1.30	1.02	1.32
FWI-EN/001: Groundwater Sampling		0.01	m	4,71		4.45	4.32	1.83
Depth		0.01	m	4./1		4.45	4.32	1.83

Page : 5 of 6
Work Order : EW1512393

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	MW10	MW11	ML-1	ML-2	ML-3
	Clie	ent sampli	ng date / time	04-Nov-2015 11:10	04-Nov-2015 11:45	04-Nov-2015 12:15	04-Nov-2015 12:00	04-Nov-2015 12:10
Compound	CAS Number	LOR	Unit	EW1512393-011	EW1512393-012	EW1512393-013	EW1512393-014	EW1512393-015
				Result	Result	Result	Result	Result
EA005FD: Field pH								
рН		0.1	pH Unit	5.6	6.1		7.1	
EA010FD: Field Conductivity								
Electrical Conductivity (Non		1	μS/cm	4570	1590		9400	
Compensated)								
EA015: Total Dissolved Solids								
Total Dissolved Solids @180°C		1	mg/L	3370	1020		6240	
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1	<1		<1	
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1	<1		<1	
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	16	44		131	
Total Alkalinity as CaCO3		1	mg/L	16	44		131	
EK055G: Ammonia as N by Discrete A	Analyser							
Ammonia as N	7664-41-7	0.01	mg/L	0.02	0.02		1.76	
EK057G: Nitrite as N by Discrete Ana	llyser							
Nitrite as N	14797-65-0	0.01	mg/L	<0.01	<0.01		0.05	
EK058G: Nitrate as N by Discrete An	alyser							
Nitrate as N	14797-55-8	0.01	mg/L	0.05	<0.01		0.05	
EK059G: Nitrite plus Nitrate as N (NC	x) by Discrete Anal	lyser						
Nitrite + Nitrate as N		0.01	mg/L	0.05	<0.01		0.10	
EK061G: Total Kjeldahl Nitrogen By D	Discrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L	2.5	2.2		3.0	
EK062G: Total Nitrogen as N (TKN + I	NOx) by Discrete An	alvser						
^ Total Nitrogen as N		0.1	mg/L	2.6	2.2		3.1	
EK067G: Total Phosphorus as P by D	iscrete Analyser							
Total Phosphorus as P		0.01	mg/L	0.14	0.07		0.05	
EN67 PK: Field Tests								
Field Observations		0.01				NO ACCESS		NO ACCESS
EP025FD: Field Dissolved Oxygen								
Dissolved Oxygen		0.01	mg/L	3.14	1.99		4.93	
FWI-EN/001: Groundwater Sampling -								1
Depth		0.01	m	2.20	2.28			
= = p = 11						<u> </u>		<u> </u>

Page : 6 of 6
Work Order : EW1512393

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Analytical Nesults								
Sub-Matrix: WATER (Matrix: WATER)		Cli	ent sample ID	ML-4	ML-5			
	CI	ient sampli	ing date / time	04-Nov-2015 12:20	04-Nov-2015 11:30			
Compound	CAS Number	LOR	Unit	EW1512393-016	EW1512393-017			
				Result	Result	Result	Result	Result
EA005FD: Field pH								
рН		0.1	pH Unit		7.1			
EA010FD: Field Conductivity								
Electrical Conductivity (Non		1	μS/cm		1820			
Compensated)								
EA015: Total Dissolved Solids								
Total Dissolved Solids @180°C		1	mg/L		1270			
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L		<1			
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L		<1			
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L		134			
Total Alkalinity as CaCO3		1	mg/L		134			
EK055G: Ammonia as N by Discrete A	nalyser							
Ammonia as N	7664-41-7	0.01	mg/L		1.65			
EK057G: Nitrite as N by Discrete Ana	lyser							
Nitrite as N	14797-65-0	0.01	mg/L		0.02			
EK058G: Nitrate as N by Discrete Ana	alyser							
Nitrate as N	14797-55-8	0.01	mg/L		0.06			
EK059G: Nitrite plus Nitrate as N (NO	x) by Discrete Ana	lyser						
Nitrite + Nitrate as N		0.01	mg/L		0.08			
EK061G: Total Kjeldahl Nitrogen By D	iscrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L		2.7			
EK062G: Total Nitrogen as N (TKN + N	NOx) by Discrete Ar	nalyser						
^ Total Nitrogen as N		0.1	mg/L		2.8			
EK067G: Total Phosphorus as P by Di	iscrete Analyser							
Total Phosphorus as P		0.01	mg/L		0.03			
EN67 PK: Field Tests								
Field Observations		0.01		NO ACCESS				
EP025FD: Field Dissolved Oxygen								
Dissolved Oxygen		0.01	mg/L		3.83			
FWI-EN/001: Groundwater Sampling -	Denth							
Depth		0.01	m					
- Arm		. •			1		l .	l .



CERTIFICATE OF ANALYSIS

Work Order : **EW1600622** Page : 1 of 6

Client : KIAMA COUNCIL Laboratory : Environmental Division NSW South Coast

Contact : MR PAUL CZULOWSKI Contact : Glenn Davies

Address : 11 MANNING STREET Address : 1/19 Ralph Black Dr, North Wollongong 2500

4/13 Geary PI, North Nowra 2541

Australia

Telephone : +61 02 4232 0444 Telephone : 02 42253125

Facsimile : +61 02 4232 0555 Facsimile : W 02 42253128 N 02 44232083

Project : Gerroa Landfill Quarterly QC Level : NEPM 2013 B3 & ALS QC Standard
Order number : 87895 Date Samples Received : 16-Feb-2016 13:38

C-O-C number : ---- Date Analysis Commenced : 16-Feb-2016

Sampler : Craig Wilson | Issue Date : 23-Feb-2016 15:46

Site : ---

KIAMA NSW, AUSTRALIA 2533

Quote number : --- No. of samples received : 17

Quote number : --- No. of samples analysed : 17

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted.

This Certificate of Analysis contains the following information:

- General Comments
- Analytical Results



NATA Accredited Laboratory 825

Accredited for compliance with ISO/IEC 17025.

Signatories

This document has been electronically signed by the authorized signatories indicated below. Electronic signing has been carried out in compliance with procedures specified in 21 CFR Part 11.

Signatories	Position	Accreditation Category
Ankit Joshi	Inorganic Chemist	Sydney Inorganics, Smithfield, NSW
Ashesh Patel	Inorganic Chemist	Sydney Inorganics, Smithfield, NSW
Dian Dao		Sydney Inorganics, Smithfield, NSW
Glenn Davies	Environmental Services Representative	Laboratory - Wollongong

Page : 2 of 6
Work Order : EW1600622

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



General Comments

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the USEPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request.

Where moisture determination has been performed, results are reported on a dry weight basis.

Where a reported less than (<) result is higher than the LOR, this may be due to primary sample extract/digestate dilution and/or insufficient sample for analysis.

Where the LOR of a reported result differs from standard LOR, this may be due to high moisture content, insufficient sample (reduced weight employed) or matrix interference.

When sampling time information is not provided by the client, sampling dates are shown without a time component. In these instances, the time component has been assumed by the laboratory for processing purposes.

Key: CAS Number = CAS registry number from database maintained by Chemical Abstracts Services. The Chemical Abstracts Service is a division of the American Chemical Society.

LOR = Limit of reporting

^ = This result is computed from individual analyte detections at or above the level of reporting

ø = ALS is not NATA accredited for these tests.

- TDS by method EA-015 may bias high for samples 5 and 8 due to the presence of fine particulate matter, which may pass through the prescribed GF/C paper.
- It has been noted that Nitrite is greater than NOx for sample 9, however this difference is within the limits of experimental variation.
- Sampling and sample data supplied by ALS Wollongong.
- Sampling completed as per FWI-EN001 Groundwater Sampling.
- Field tests completed on day of sampling/receipt.

Page : 3 of 6
Work Order : EW1600622

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	MW1D	MW1S	MW3	MW4	MW5	
	Cli	ent sampli	ng date / time	16-Feb-2016 10:21	16-Feb-2016 10:20	16-Feb-2016 10:11	16-Feb-2016 10:34	16-Feb-2016 10:03	
Compound	CAS Number	LOR	Unit	EW1600622-001	EW1600622-002	EW1600622-003	EW1600622-004	EW1600622-005	
				Result	Result	Result	Result	Result	
EA005FD: Field pH									
рН		0.1	pH Unit	7.3		7.0	6.1	7.0	
EA010FD: Field Conductivity									
Electrical Conductivity (Non		1	μS/cm	631		822	207	123	
Compensated)									
EA015: Total Dissolved Solids									
Total Dissolved Solids @180°C		1	mg/L	419		476	226	162	
ED037P: Alkalinity by PC Titrator									
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1		<1	<1	<1	
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1		<1	<1	<1	
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	252		137	72		
Total Alkalinity as CaCO3		1	mg/L	252		230	137	72	
EK055G: Ammonia as N by Discrete	Analyser								
Ammonia as N	7664-41-7	0.01	mg/L	2.43		20.9	0.12	<0.01	
EK057G: Nitrite as N by Discrete Ana	alyser								
Nitrite as N	14797-65-0	0.01	mg/L	0.01		0.02	<0.01	<0.01	
EK058G: Nitrate as N by Discrete An	alvser								
Nitrate as N	14797-55-8	0.01	mg/L	0.02		0.02	0.02	0.02	
EK059G: Nitrite plus Nitrate as N (NC	Dx) by Discrete Anal	lvser							
Nitrite + Nitrate as N		0.01	mg/L	0.03		0.04	0.02	0.02	
EK061G: Total Kjeldahl Nitrogen By D	Discrete Analyser								
Total Kjeldahl Nitrogen as N		0.1	mg/L	3.3		58.1	0.6	1.0	
EK062G: Total Nitrogen as N (TKN + I	NOx) by Discrete An	alvser	-						
^ Total Nitrogen as N		0.1	mg/L	3.3		58.1	0.6	1.0	
EK067G: Total Phosphorus as P by D	Discrete Analyser		3						
Total Phosphorus as P		0.01	mg/L	0.17		2.71	1.08	0,26	
EN67 PK: Field Tests			J. =					V	
Field Observations		0.01			DRY				
EP025FD: Field Dissolved Oxygen									
Dissolved Oxygen		0.01	mg/L	3,70		2.80	2.40	3.20	
		0.01	mg/L	5.70		2.00	2.70	J.20	
FWI-EN/001: Groundwater Sampling - Depth		0.01	m	3.60		4.10	4.40	3.90	
Бериі		0.01	111	3.00		4.10	4.40	3.90	

Page : 4 of 6
Work Order : EW1600622

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	MW6D	MW6S	MW7D	MW7S	MW9	
	Cli	ient sampli	ng date / time	16-Feb-2016 09:23	16-Feb-2016 09:20	16-Feb-2016 09:54	16-Feb-2016 09:44	16-Feb-2016 11:04	
Compound	CAS Number	LOR	Unit	EW1600622-006	EW1600622-007	EW1600622-008	EW1600622-009	EW1600622-010	
				Result	Result	Result	Result	Result	
EA005FD: Field pH									
рН		0.1	pH Unit	6.8		7.3	7.3	6.3	
EA010FD: Field Conductivity									
Electrical Conductivity (Non		1	μS/cm	732		443	579	9520	
Compensated)									
EA015: Total Dissolved Solids									
Total Dissolved Solids @180°C		1	mg/L	486		382	434	6840	
ED037P: Alkalinity by PC Titrator									
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1		<1	<1	<1	
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1		<1	<1	<1	
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	318		170	84		
Total Alkalinity as CaCO3		1	mg/L	318		182	170	84	
EK055G: Ammonia as N by Discrete	Analyser								
Ammonia as N	7664-41-7	0.01	mg/L	3.48		1.02	0.70	0.07	
EK057G: Nitrite as N by Discrete Ana	alyser								
Nitrite as N	14797-65-0	0.01	mg/L	<0.01		<0.01	0.02	<0.01	
EK058G: Nitrate as N by Discrete An	alvser								
Nitrate as N	14797-55-8	0.01	mg/L	0.26		0.03	<0.01	0.02	
EK059G: Nitrite plus Nitrate as N (NC	(Dx) by Discrete Ana	lvser							
Nitrite + Nitrate as N		0.01	mg/L	0.26		0.03	<0.01	0.02	
EK061G: Total Kjeldahl Nitrogen By [Discrete Analyser								
Total Kjeldahl Nitrogen as N		0.1	mg/L	4.2		1.4	1.2	1.4	
EK062G: Total Nitrogen as N (TKN + I	NOx) by Discrete An	alvser	-						
^ Total Nitrogen as N		0.1	mg/L	4.5		1.4	1.2	1.4	
EK067G: Total Phosphorus as P by D	liscroto Analysor		3						
Total Phosphorus as P		0.01	mg/L	4.55		1.60	0.32	0.03	
EN67 PK: Field Tests			J. =					*****	
Field Observations		0.01			DRY				
EP025FD: Field Dissolved Oxygen		J.J.							
Dissolved Oxygen		0.01	mg/L	2.80		3.10	3.90	1.90	
		0.01	mg/L	2.00		3.10	0.30	1.30	
FWI-EN/001: Groundwater Sampling - Depth		0.01	m	5.00		4.80	4.70	1.80	
Бериі		0.01	111	5.00		4.00	4.70	1.00	

Page : 5 of 6
Work Order : EW1600622

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	MW10	MW11	ML-1	ML-2	ML-3	
	Cli	ient sampli	ng date / time	16-Feb-2016 10:51	16-Feb-2016 11:17	16-Feb-2016 11:50	16-Feb-2016 11:32	16-Feb-2016 11:53	
Compound	CAS Number	LOR	Unit	EW1600622-011	EW1600622-012	EW1600622-013	EW1600622-014	EW1600622-015	
			Result	Result	Result	Result	Result		
EA005FD: Field pH									
pH		0.1	pH Unit	5.1	5.6		6.9		
EA010FD: Field Conductivity									
Electrical Conductivity (Non		1	μS/cm	8060	6370		25400		
Compensated)									
EA015: Total Dissolved Solids									
Total Dissolved Solids @180°C		1	mg/L	6310	4510		17200		
ED037P: Alkalinity by PC Titrator									
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L	<1	<1		<1		
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L	<1	<1		<1		
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L	8	34	102			
Total Alkalinity as CaCO3		1	mg/L	8	34		102		
EK055G: Ammonia as N by Discrete A	Analyser								
Ammonia as N	7664-41-7	0.01	mg/L	<0.01	<0.01		0.81		
EK057G: Nitrite as N by Discrete Ana	llyser								
Nitrite as N	14797-65-0	0.01	mg/L	<0.01	<0.01		0.02		
EK058G: Nitrate as N by Discrete Ana	alyser								
Nitrate as N	14797-55-8	0.01	mg/L	0.09	<0.01		0.02		
EK059G: Nitrite plus Nitrate as N (NO	x) by Discrete Ana	lyser							
Nitrite + Nitrate as N		0.01	mg/L	0.09	<0.01		0.04		
EK061G: Total Kjeldahl Nitrogen By D	iscrete Analyser								
Total Kjeldahl Nitrogen as N		0.1	mg/L	1.7	5.0		1.6		
EK062G: Total Nitrogen as N (TKN + N	NOx) by Discrete An	alvser_						•	
^ Total Nitrogen as N		0.1	mg/L	1.8	5.0		1.6		
EK067G: Total Phosphorus as P by D	iscrete Analyser		_						
Total Phosphorus as P		0.01	mg/L	0.06	0.26		0.03		
EN67 PK: Field Tests			<u> </u>						
Field Observations		0.01				NO ACCESS		NO ACCESS	
EP025FD: Field Dissolved Oxygen								1 11	
Dissolved Oxygen		0.01	mg/L	2.90	2.00		5.40		
		0.01	g, _	2.00	2.00		0.40		
FWI-EN/001: Groundwater Sampling - Depth		0.01	m	2.10	2.30				
Берин		0.01	111	2.10	2.30				

Page : 6 of 6
Work Order : EW1600622

Client : KIAMA COUNCIL
Project : Gerroa Landfill Quarterly



Sub-Matrix: WATER (Matrix: WATER)		Clie	ent sample ID	ML-4	ML-5			
	Cli	ent sampli	ng date / time	16-Feb-2016 11:55	16-Feb-2016 11:06			
Compound	CAS Number	CAS Number LOR Unit		EW1600622-016	EW1600622-017			
				Result	Result	Result	Result	Result
EA005FD: Field pH								
рН		0.1	pH Unit		6.8			
EA010FD: Field Conductivity								
Electrical Conductivity (Non		1	μS/cm		9900			
Compensated)								
EA015: Total Dissolved Solids								
Total Dissolved Solids @180°C		1	mg/L		6860			
ED037P: Alkalinity by PC Titrator								
Hydroxide Alkalinity as CaCO3	DMO-210-001	1	mg/L		<1			
Carbonate Alkalinity as CaCO3	3812-32-6	1	mg/L		<1			
Bicarbonate Alkalinity as CaCO3	71-52-3	1	mg/L		102			
Total Alkalinity as CaCO3		1	mg/L		102			
EK055G: Ammonia as N by Discrete A	nalyser							
Ammonia as N	7664-41-7	0.01	mg/L		1.24			
EK057G: Nitrite as N by Discrete Ana	lyser							
Nitrite as N	14797-65-0	0.01	mg/L		0.02			
EK058G: Nitrate as N by Discrete Ana	alyser							
Nitrate as N	14797-55-8	0.01	mg/L		<0.01			
EK059G: Nitrite plus Nitrate as N (NO	x) by Discrete Anal	lyser						
Nitrite + Nitrate as N		0.01	mg/L		0.02			
EK061G: Total Kjeldahl Nitrogen By D	iscrete Analyser							
Total Kjeldahl Nitrogen as N		0.1	mg/L		2.5			
EK062G: Total Nitrogen as N (TKN + N	IOx) by Discrete An	alyser						
^ Total Nitrogen as N		0.1	mg/L		2.5			
EK067G: Total Phosphorus as P by Di	iscrete Analyser							
Total Phosphorus as P		0.01	mg/L		0.03			
EN67 PK: Field Tests								
Field Observations		0.01		NO ACCESS				
EP025FD: Field Dissolved Oxygen								
Dissolved Oxygen		0.01	mg/L		5.10			
FWI-EN/001: Groundwater Sampling -								1
Depth		0.01	m					
							<u> </u>	<u> </u>



Appendix B

					Toolijooa	a (rainfall 2	2015)						Bombo	(rainfall)
2015	Jan-15	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec-15	Jan-16	Feb-16
Graph	ilit	ilii	ilit	ilit	thi	ilit	ilit	thi	ilii	ilit	thi	thi	ilit	thi
1st	0	0	0	9.4	22.8	0.4	0	0	0	1.2	0.4	0	0	0
2nd	0	1.6	8.2	0	16.2	0	0	0	0	0	0.6	0	0	13.8
3rd	0	0	0	3.6	11.2	0	0	0	6.2	0	17.6	0	0	0
4th	0	0.6	0	11	1.2	0	0	0	10.6	0	20	0	10.6	6.2
5th	35.8	7.8	0	13.8	0.4	2.6	0	0	0	0	23.8	0	11.8	0.2
6th	0	0	0	0	0	0	0	0	0	0	14.4	0	10	0
7th	0	0	0	17.6	0.2	0	0	0	0	0	2.8	0	21.2	0
8th	0	0	0	9.4	0	0	3	0	0	0	4.4	1.6		0
9th	1.4	0	0	0.8	1.4	0.8	1.2	0	0	0	0.6	0	0	0
10th	0	14.4	0	0	0	0	0	0	0	0.8	0	0	0	0
11th	53.4	3.2	0	28	0	0	0	0	0	0	0	0	0	0
12th	23	0	0	2	0	0	8.8	0	0	0	5.4	0	0	0
13th	6.2	1	8.4	1	0	0	2.6	0	0	3.8	1	0	0	0
14th	0	2	0	0	0	0	0	0	0	3.2	1.8	0	0	0
15th	0	22.4	0	0	0	0	0	0	0	0	12.2	0	24.6	0
16th	0.4	0	5.2	0	0	26.8	0	0	0	0	0.6	0.4	1.4	0
17th	0	0	0	5.6	0	1	38.2	0	0.8	0	0	11.8	0.2	2.4
18th						5.22								
10111	0	0	0	2.2	0	days	1.4	1.2	0.4	1.8	0	0	0	0
19th	22.6	1.2	0	8	0	47.2	0	0	16.2	0	0	0	0	0
20th	13.2	0	0	43	0	13	0	0	0	0	0	0	0	0
21st	14.2	1.6	0	68.8	0	0.8	0	0	0	0	0	0	0	1.4
22nd	0	4.2	6.4	108.4	21.4	0	0	0	0	16.8	1	31	11.4	0
23rd	0	0	0	71.6	24.6	0	0	0.6	0	12.8	0	14.4	11.2	0
24th	0	0	0	0	0	0	0	5.6	4.8	0	0	0.2	0.6	0
25th	0.2	48.4	37	0	0	2	0	174.4	7	0	0	0	0	0
26th	0.2	3.8	0	11.8	0	0.6	0	177	15	0	0	0	10.8	0
27th	53.2	0.2	0	4.4	0	0.4	0	10	3.2	26.8	0	0	0	1.2
28th	85	0	0	0	0	6	0	0	0	0.6	0	82	3.6	0.4
29th	7		0	0	0.6	0	0	9.2	0	0	0	0	0	0
30th	0		0	6	0	0	0	0	0	0	0.2	0	5.8	
31st	0.2		0		0		0	0		0		0	1.4	
Highest														
Daily	85	48.4	37	108.4	24.6	47.2	38.2	177	16.2	26.8	23.8	82	24.6	13.8
Monthly Total	316	112.4	65.2	426.4	100	105.8	55.2	378	64.2	67.8	106.8	141.4	124.6	25.6

of daily data

Feb 2015 to end Jan 2016 17	47.8
-----------------------------	------




Appendix C: Ammonia Trigger Values

Appendix C

Ammonia levels and 20% trigger factor for Groundwater and 10% trigger factor for Surface water at Gerroa Landfill

Date Sampled	MW1 D	MW1 S	MW3	MW4	MW5	MW6 D	MW6 S	MW7 D	MW7 S	MW9	MW10	MW11	ML-1	ML-2	ML-3	ML-	ML-5
20% trigger level (mg/L)	164.4	1.32	39.24	21.8	7.07	98.8	27.78	85.32	14.3	4.26	2.85	2.96					
10% trigger level (mg/L)													0.836	1.38 Exceed Nov 2015 #1.76 mg/L)	0.066	0.23	2.38 Exceed Aug 2015 #2.39 mg/L)

Legend:



Feedback is welcomed.

Dino Parisotto (dino@earth2water.com.au)

